


Estia Eichten
Fermilab

- The Long View
- Return to the Energy Frontier
- Staging Physics Milestones
- Summary



2013 Community Summer Study
"Snowmass on the Mississippi"
University of Minnesota
July 29-Aug 6, 2013



WhitePapers

1. Enabling Intensity and Energy Frontier Science with a Muon Accelerator Facility In the USA
2. Muon Collider Higgs Factory


Three Pillars of nuSTORM



- Delivers on the physics for the study of sterile ν
 - Offering a new approach to the production of ν beams setting a 10σ benchmark to make definitive statement w/r LSND/MiniBooNE
- Can add significantly to our
- ★ knowledge of ν interactions, particularly for ν_e
 - ν "Light Source"
- ★ Provides an accelerator & detector technology test bed

Which Accelerator for Higgs Physics?



1. The LHC is the Higgs Accelerator - Continue -> HL-LHC
2. Continue research and development of lepton colliders. In particular the muon collider needs a convincing proof of 6D cooling.
3. Push neutrino physics - Lepton sector
4. After 300 fb^{-1} of $\sim 14 \text{ TeV}$ running OR the discovery of BSM physics, chose the next accelerator for Higgs physics.








Implications of early LHC Results

- No evidence for new physics beyond the Standard Model (BSM) to date:
 - BSM (SUSY, Strong Dynamics, Extra Dimensions, New fermions or gauge bosons,...)
 - ATLAS limits



CMS limits

CMS EXOTICA 95% CL EXCLUSION LIMITS (TeV)



- Scales already probed at the LHC suggest that to study BSM new physics the next energy frontier collider must have \sqrt{s} in the multi-TeV range even for EW processes.
- However there must be new physics !!! WHY? Let me list the reasons

1. The Standard Model is incomplete:

- dark matter; neutrino masses and mixing \rightarrow new fields or interactions;
- baryon asymmetry in the universe \rightarrow more CP violation
- gauge unification \rightarrow new interactions;
- gravity: strings and extra dimensions

2. Experimental hints of new physics: $(g-2)_\mu$, top A_{fb} , ...

3. Theoretical problems with the SM:

- Scalar sector problematic:


$$\mu^2 (\Phi^\dagger \Phi) + \lambda (\Phi^\dagger \Phi)^2 + \Gamma_{ij} \psi_{iL}^\dagger \psi_{jR} \Phi + \text{h.c.}$$

$m_H^2/M_{\text{planck}}^2 \approx 10^{-34}$
Hierarchy problem

vacuum
stability

large range of
fermion masses


- The SM Higgs boson is unnatural. (m_H^2/μ^2)
- Solutions: SUSY, New Strong Dynamics, ...



muon $(g-2)$

Davier, Hoecker,
Malaescu, Zhang
Jegerlehner, Szafron
Hagiwara, Liao,
Martin, Nomura,
Teubner

hadronic VP
contributions
 $(685 \pm 4) \times 10^{-10}$






There remains a persistent discrepancy of 3.3-3.6 σ



- The strong case for a TeV scale hadron collider rested on two arguments:
 1. Unitarity required that a mechanism for EWSB was manifest at or below the TeV scale.
 2. The SM is unnatural ('t Hooft conditions) and incomplete (dark matter, insufficient CP violation for the observed baryon excess, gauge unification, gravity and strings)
- If after the analysis of the 2012 CMS/ATLAS data, the 126 GeV state is found to be a 0^+ state with couplings consistent with the SM Higgs, the first argument is satisfied.
 - The second argument remains strong. but is less strongly tied to the TeV scale.
 - Scales already probed at the LHC suggest that any new collider (of LHC level costs) should be able to probe the BSM physics in the multi-TeV range.

Muon Collider

- $\mu^+ \mu^-$ Collider:
 - Center of Mass energy: 1.5 - 6 TeV (3 TeV)
 - Luminosity $> 10^{34} \text{ cm}^{-2} \text{ sec}^{-1}$ (440 fb $^{-1}$ /yr)
 - Compact facility
 - 3 TeV - ring circumference 3.8 km
 - 2 Detectors
 - Superb Energy Resolution
 - MC: 95% luminosity in $dE/E \sim 0.1\%$
 - CLIC: 35% luminosity in $dE/E \sim 1\%$






- Comparison of Lepton Colliders at High Energy




- For $\sqrt{s} < 500 \text{ GeV}$
 - SM thresholds: $Z^0 h$, $W^+ W^-$, top pairs
 - Higgs factory ($\sqrt{s} \approx 126 \text{ GeV}$) ✓
- For $\sqrt{s} > 500 \text{ GeV}$
 - Sensitive to possible Beyond SM physics.
 - High luminosity required. ✓
 - Cross sections for central ($|\theta| > 10^\circ$) pair production $\sim R \times 86.8 \text{ fb/s (in TeV}^2\text{)} (R \approx 1)$
 - At $\sqrt{s} = 3 \text{ TeV}$ for $100 \text{ fb}^{-1} \sim 1000 \text{ events/(unit of R)}$
- For $\sqrt{s} > 1 \text{ TeV}$
 - Fusion processes important at multi-TeV MC

$$\sigma(s) = C \ln\left(\frac{s}{M_X^2}\right) + \dots$$

- An Electroweak Boson Collider ✓



- But muons decay:
 - The muon beams must be accelerated and cooled in phase space (factor $\approx 10^6$) rapidly
→ ionization cooling
 - requires a complex cooling scheme
 - The decay products ($\mu^- \rightarrow \nu_\mu \nu_e e^-$) have high energies.
 - Detector background issues
 - Neutrino beam issue for $E_{cm} \geq 4$ TeV. Beam steering resolves this for $E_{cm} \lesssim 10$ TeV.
- The issues need dedicated R&D
 - MICE
 - MAP
 - nuStorm - Definitive 6D cooling demo.



Staging A Muon Collider

- Provides a flexible staging scenerio with physics at each stage.

- Neutrino Factory




Figure 15: Functional elements of a 5 GeV Neutrino Factory

- Higgs Factory
- High Energy Muon Collider




Figure 26: Functional elements of a Higgs Factory/Muon Collider complex

Staging Scenerio


- A possible timeline

- Project X Stages:

- Stage I -> 1 GeV, 1 mA
- Stage II -> 3 GeV, 3MW
- Stage III -> 8 GeV
- Stage IV -> 4MW

- Decision points:

- Finish of MAP Feasibility Assessment ~ 2018
- Advanced System R&D makes use of nuSTORM muon ring.
- Decision point middle of 2020's on collider program.
- Program X Stage II can start physics of neutrino or collider program.



• Neutrino Physics Staging

- Because θ_{13} is large a lower energy (5 GeV) and 1300 km works for a Neutrino Factory.
- First a lower intensity (Project X phase 2) ($2 \times 10^{20} \mu^\pm/\text{yr}$) neutrino factory NuMAX
- Then higher intensity ($1.2 \times 10^{21} \mu^\pm/\text{yr}$) NuMAX+
- Unsurpassed performance is obtained for 34 kton magnetized LAr (TPC) at distance 1300 km (NuMAX+)




Table 1. Muon Accelerator Program baseline Neutrino Factory parameters for nuSTORM and two NuMAX phases located on the Fermilab site and pointed towards a detector at SURF. For comparison, the parameters of the IDS-NF are also shown.

System	Parameters	Unit	nuSTORM	NuMAX	NuMAX+	IDS-NF
Performance	Stored μ^\pm or μ^-/year		8×10^{17}	2×10^{20}	1.2×10^{21}	1×10^{21}
	ν_e or ν_μ to detectors/yr		3×10^{17}	8×10^{19}	5×10^{20}	5×10^{20}
Detector	Far Detector:	Type	SuperBIND	MIND / Mag LAr	MIND / Mag LAr	MIND
	Distance from Ring	km	1.9	1300	1300	2000
	Mass	kT	1.3	30 / 10	100 / 30	100
	Magnetic Field	T	2	0.5-2	0.5-2	1-2
	Near Detector:	Type	SuperBIND	Suite	Suite	Suite
	Distance from Ring	m	50	100	100	100
Neutrino Ring	Mass	kT	0.1	1	2.7	2.7
	Magnetic Field	T	Yes	Yes	Yes	Yes
	Ring Momentum (P_μ)	GeV/c	3.8	5	5	10
	Circumference (C)	m	480	600	600	1190
	Straight section	m	185	235	235	470
	Arc Length	m	50	65	65	125
Acceleration	Initial Momentum	GeV/c	-	0.22	0.22	0.22
	Single-pass Linac	GeV/pass	-	0.95	0.95	0.56
		MHz	-	325	325	201
	4.5-pass RLA	RLA I	GeV/pass	0.85	0.85	0.45
		RLA I	MHz	325	325	201
		RLA II	GeV/pass	-	-	1.6
		RLA II	MHz	-	-	201
			-	-	-	-
Cooling			No	No	4D	4D
Proton Source	Proton Beam Power	MW	0.2	1	3	4
	Proton Beam Energy	GeV	120	3	3	10
	Protons/year	1×10^{21}	0.1	41	125	25
	Repetition Frequency	Hz	0.75	70	70	50

- Staging Steps:

- Higgs factory $\sqrt{s} = m_H \approx 126 \text{ GeV}$

- $\mathcal{L} = 1.7 \times 10^{31} \sim 170 \text{ pb}^{-1} / \text{yr};$
 $\Delta E/E = 0.003\%$
- $\mathcal{L} = 8 \times 10^{31} \sim 800 \text{ pb}^{-1} / \text{yr};$
 $\Delta E/E = 0.004\%$

- High Energy Muon Collider:

- LHC at $\sqrt{s} \approx 14 \text{ TeV}$ after 300 fb^{-1} .
Muon collider design energy is flexible. ($\Delta E/E = 0.1\%$)
- $\sqrt{s} = 1.5 \text{ TeV};$
 $\mathcal{L} = 1.25 \times 10^{34} \sim 125 \text{ fb}^{-1} / \text{yr};$
- $\sqrt{s} = 3.0 \text{ TeV};$
 $\mathcal{L} = 4.4 \times 10^{34} \sim 440 \text{ fb}^{-1} / \text{yr}$
- $\sqrt{s} = 6.0 \text{ TeV};$
 $\mathcal{L} = 1.6 \times 10^{35} \sim 1.6 \text{ ab}^{-1} / \text{yr}$

Muon Collider Baseline Parameters					
Parameter	Units	Higgs Factory		Multi-TeV Baselines	
		Startup Operation	Production Operation		
CoM Energy	TeV	0.126	0.126	1.5	3.0
Avg. Luminosity	$10^{34} \text{ cm}^{-2} \text{ s}^{-1}$	0.0017	0.008	1.25	4.4
Beam Energy Spread	%	0.003	0.004	0.1	0.1
Higgs/ 10^7 sec		3,500	13,500	37,500	200,000
Circumference	km	0.3	0.3	2.5	4.5
No. of IPs		1	1	2	2
Repetition Rate	Hz	30	15	15	12
β^*	cm	3.3	1.7	1 (0.5-2)	0.5 (0.3-3)
No. muons/bunch	10^{12}	2	4	2	2
No. bunches/beam		1	1	1	1
Norm. Trans. Emittance, ϵ_{TN}	$\pi \text{ mm-rad}$	0.4	0.2	0.025	0.025
Norm. Long. Emittance, ϵ_{LN}	$\pi \text{ mm-rad}$	1	1.5	70	70
Bunch Length, σ_s	cm	5.6	6.3	1	0.5
Beam Size @ IP	μm	150	75	6	3
Beam-beam Parameter / IP		0.005	0.02	0.09	0.09
Proton Driver Power	MW	4 [#]	4	4	4

[#] Could begin operation with Project X Stage 2 beam

- A muon collider can directly produce the Higgs as an s-channel resonance.

- Higgs couples to mass so rate enhanced by $\left[\frac{m_\mu}{m_e}\right]^2 = 4.28 \times 10^4$ so the cross section is $\sigma(\mu^+\mu^- \rightarrow h) = 26 \text{ pb}$ (for $\Delta = \Gamma$ and including ISR and a 15° forward cut).
- To obtain the same sensitivity to Higgs decay modes in a electron collider via Zh process as s-channel production at a MC requires more than 100 times the integrated luminosity.
- The excellent energy resolution Δ of a muon collider makes the process observable.

$$\sigma_{\text{eff}}(s) = \int d\sqrt{\hat{s}} \frac{dL(\sqrt{\hat{s}})}{d\sqrt{\hat{s}}} \sigma(\mu^+\mu^- \rightarrow h \rightarrow X)$$

$$\propto \begin{cases} \Gamma_h^2 B / [(s - m_h^2)^2 + \Gamma_h^2 m_h^2] & (\Delta \ll \Gamma_h), \\ B \exp\left[-\frac{(m_h - \sqrt{s})^2}{2\Delta^2}\right] (\frac{\Gamma_h}{\Delta}) / m_h^2 & (\Delta \gg \Gamma_h). \end{cases}$$


$$\sigma(\mu^+\mu^- \rightarrow h \rightarrow X) = \frac{4\pi\Gamma_h^2 \text{Br}(h \rightarrow \mu^+\mu^-) \text{Br}(h \rightarrow X)}{(\hat{s} - m_h^2)^2 + \Gamma_h^2 m_h^2}.$$

- Results:

Channel	δM_H (MeV)	$\delta \Gamma_H$ (MeV)	$\delta \text{Br}(h \rightarrow X)$
$b\bar{b}$	0.1	0.4	0.05
WW^*	0.07	0.2	0.01
Combined	0.06	0.18	—

- $\Delta \text{Br}(\mu^+\mu^-) \text{Br}(WW^*) \sim 2\%$

- Finding the Higgs (5σ) requires 270 pb^{-1} .




- New Z' , W'
 - S-channel resonances - factories for lepton colliders
 - Set minimum luminosity for MC.

Minimum luminosity at Z' peak:

$$\mathcal{L} = 1.0\text{--}5.0 \times 10^{30} \text{ cm}^{-2} \text{ sec}^{-1}$$

for $M(Z') \rightarrow 2.5\text{--}5.0 \text{ TeV}$



- A muon collider can be built to operate well above 4 TeV :

- Keeping the same limits on neutrino radiation.

The luminosity will scale as:

$$L(E_{cm})/L(4 \text{ TeV}) = [E_{cm}/(4 \text{ TeV})]^{-2}$$

- If the emittance can be reduced as the energy is increased, up to one power of energy ratio can be recovered.

- Hence an s-channel resonance well in excess of 10 TeV could be studied in detail at such a muon collider.

- Two Higgs doublets (MSSM):



- Five scalar particles: h^0, H^0, A^0, H^\pm
- Decay amplitudes depend on two parameters: (α, β)

	$\mu^+\mu^-, b\bar{b}$	$t\bar{t}$	ZZ, W^+W^-	ZA^0
h^0	$-\sin\alpha/\cos\beta$	$\cos\alpha/\sin\beta$	$\sin(\beta-\alpha)$	$\cos(\beta-\alpha)$
H^0	$\cos\alpha/\cos\beta$	$\sin\alpha/\sin\beta$	$\cos(\beta-\alpha)$	$-\sin(\beta-\alpha)$
A^0	$-i\gamma_5 \tan\beta$	$-i\gamma_5/\tan\beta$	0	0


$$\tan 2\alpha = \frac{M_A^2 + M_Z^2}{M_A^2 - M_Z^2} \tan 2\beta.$$

- decoupling limit $m_{A^0} \gg m_{Z^0}$:

- » h^0 couplings close to SM values
- » H^0, H^\pm and A^0 nearly degenerate in mass
- » H^0 small couplings to VV , large couplings to ZA^0
- » For large $\tan\beta$, H^0 and A^0 couplings to charged leptons and bottom quarks enhanced by $\tan\beta$. Couplings to top quarks suppressed by $1/\tan\beta$ factor.




- The LHC has difficulty observing the H, A especially for masses > 500 GeV. Even at $\sqrt{s} = 14$ TeV and 300 fb^{-1} .



Significance contours for SUSY Higgses

Regions of the MSSM parameter space ($m_A, \tan\beta$) explorable through various SUSY Higgs channels

- 5σ significance contours
- two-loop / RGE-improved radiative corrections
- $m_{\text{top}} = 175 \text{ GeV}$, $m_{\text{SUSY}} = 1 \text{ TeV}$, no stop mixing ;




- Pair produced with easy at a multi-TeV lepton collider.

- If H/A near present LHC bounds. The states can be cleanly separated because of the excellent energy resolution of the muon collider.



- Example of Natural SUSY

- Low-lying spectrum




- For electroweakinos, sleptons, ...

A ≥ 3 TeV muon collider has discovery reach beyond a 100 TeV pp collider !



Dittmaier and Kaiser
[hep-ph/0203120]

- Generally expect very heavy: H^\pm , H^0 and A^0
 - LHC limits on H^\pm : ~ 300 (ATLAS) (CMS)
 - SUSY models that evade the all present experimental constraints often have very heavy THDM scalars
- The H/A are observable as s-channel resonances at a MC!
 - $M_H = M_A \sim 1.5 \text{ TeV}/c^2$, $\Gamma \sim 15 \text{ GeV}$
 - Large $\tan\beta \sim 20$
 - Limited spectrum of SUSY particle decays.
 - Expect 10^6 H/A decays per 1 ab^{-1}
- The H/A resonances are a factory for study BSM physics.
 - E.E and A. Martin (arXiv:1306.2609)



- Electroweak Symmetry Breaking is generated dynamically at nearby scale
 - Technicolor, ETC, walking TC, topcolor, Two Scale TC, composite Higgs models, ...
 - New strong interaction at the Terascale:
 - What is the spectrum of low-lying states? s -channel production π_T (technipion) (0^-), ρ_T , ω_T (technirho, techniomega) nearly degenerate - needs good energy resolution
 - What is the ultraviolet completion? Gauge group? Fermion representations?
 - What is the energy scale of the new dynamics?
 - Any new insight into quark and/or lepton flavor mixing and CP violation?


- Contact interactions

- e.g. Compositeness, broken flavor symmetries, ...

- Present LHC bounds (~ 10 TeV)
$$\mathcal{L} = \frac{g^2}{\Lambda^2} (\bar{\Psi} \Gamma \Psi) (\bar{\Psi} \Gamma' \Psi)$$

- Muon collider sensitive to scales > 200 TeV

- Forward cone cut not important
- Polarization useful in determining chiral character of the interaction.



- The observation of a new state at 125 GeV by both ATLAS and CMS revitalizes consideration of a Higgs factory as part of a staged multi-Tev muon collider. This is particularly attractive if there is an enlarged scalar sector (eg. THDM, SUSY)
- The unique measurements of the Muon Higgs factory (4.2 fb^{-1})
 - Most precise measurement of Higgs mass: $\Delta m_H = 0.06 \text{ MeV}$
 - Direct Higgs width measurement: $\Delta \Gamma_H = 0.18 \text{ MeV}$.
 - Measurement of $\text{BR}(\mu^+\mu^-) \text{BR}(WW^*)$ to 2%.
 - Disentangle nearly degenerate scalar resonances.
- A multiTeV lepton collider will be required for full coverage of Terascale physics.
 - The physics potential for a muon collider at $\sqrt{s} \sim 3 \text{ TeV}$ and integrated luminosity of 1 ab^{-1} is outstanding. Particularly strong case for SUSY and new strong dynamics.
 - Narrow s-channel states played an important role in past lepton colliders. If such states exist in the multi-TeV region, they will play a similar role in precision studies for new physics.
- A staged Muon Collider can also provide a Neutrino Factory to fully disentangle neutrino physics.
- It all starts with nuStorm !!

BACKUP SLIDES

- Concept of naturalness.
 - K. Wilson, G. 't Hooft
 - A theory $[L(\mu)]$ is natural at scale $\mu \Leftrightarrow$ for any small dimensionless parameter λ (e.g. m/μ) in $L(\mu)$ the limit $\lambda \rightarrow 0$ enhances the symmetries of $L(\mu)$
- The SM Higgs boson is unnatural. (m_H^2/μ^2)
 - Maybe no large gap in scales (Extra Dimensions)
- Two potential solutions:
 - scalars not elementary
 - New strong dynamics (TC, walking TC, little Higgs, top color, ...)
 - fermion masses are natural
 - Symmetry coupling fermions and bosons (SUSY)
- Quest for the "natural" theory to replace the SM has preoccupied theorists since the early 80's
- Is a third way required after the discovery of a Higgs boson?

G. 't Hooft in Proceedings of
Recent Developments in Gauge Theories,
Cargese, France (1980)

NATURALNESS, CHIRAL SYMMETRY, AND SPONTANEOUS

CHIRAL SYMMETRY BREAKING

G. 't Hooft

Institute for Theoretical Physics

Utrecht, The Netherlands

ABSTRACT

A properly called "naturalness" is imposed on gauge theories. It is an order-of-magnitude restriction that must hold at all energy scales μ . To construct models with complete naturalness for elementary particles one needs more types of confining gauge theories besides quantum chromodynamics. We propose a search