

Constraining the top-Z coupling through $t\bar{t}Z$ production at the LHC

Raoul Röntschi



Loopfest XIII
18 June 2014

RR and Markus Schulze
arXiv:hep-ph/1404.1005

- Motivation
 - Why top physics is interesting
 - Why NLO calculation with spin-correlated decays
- Top-Z Lagrangian in effective field theory
- Details of calculation
- Constraints from current CMS data
- Constraints from future LHC run
- Conclusions and future work


- Top quark has **largest mass** of known quarks.
- Decays before hadronization.
- $y_t \sim 1$.
- Special role in EWSB?
- Expect top-EW couplings to be **highly sensitive** to EWSB mechanism.

⇒ measurement of top-EW couplings is **part of the program of understanding EWSB**, as well as avenue for **finding/bounding New Physics effects**.

Top quarks at the LHC

LHC a **top factory**: $\sigma_{t\bar{t}} \sim nb$ at $\sqrt{s} = 14$ TeV

$\Rightarrow \sim 10^9$ top pairs over lifetime of LHC.



Campbell, Ellis, RR, hep-ph/1302.3856

- $t\bar{t} + \gamma$, $t\bar{t}Z$ and $t\bar{t}W$ observed in run I.

hep-ex/1307.4568, ATLAS-CONF-2012-126, hep-ex/1303.3239

- Low statistics in $t\bar{t}Z$ channel: CMS 9 events, ATLAS 1 event.

Long-term project requiring high luminosity at higher energy run.

Energy and luminosity large enough to produce **massive EW particles** in association with $t\bar{t}$.

\rightarrow **direct measurement of top-EW couplings**

No **direct constraints** placed on top-Z couplings to date.

Indirect constraints through LEP data:

- Zbb coupling and ρ parameter closely constrained by fits to ϵ_1 and ϵ_b .
- R_b and A_{FB}^b also constrain $Zb_L b_L$ couplings \rightarrow constrain $Zt_L t_L$ coupling (under assumption of $SU(2)$ symmetry).
- Translate into $\sim 1\%$ constraints on top-Z coupling.

Unlikely that LHC will be able to improve on this.

COMPLEMENT, not COMPETE.

How well can the top-Z coupling be constrained at the $\sqrt{s} = 13$ TeV LHC?

- Luminosity – e.g. 30, 300, 3000 fb⁻¹
- Experimental accuracy
- Theoretical accuracy

Cf. Baur, Juste, Orr, Rainwater:

hep-ph/0412021, hep-ph/0512262

- 1 Trileptonic channel $t\bar{t}Z \rightarrow (jjbb\nu l^+ l^-)$ best compromise between clean signal and overall rate.
- 2 Shape of opening angle between leptons from Z decay $\Delta\phi_{ll}$ sensitive to top-Z couplings.
- 3 Scale uncertainty is biggest obstacle (on theoretical side)!

Motivated by this, we perform a **partonic level** calculation to **NLO** in pQCD.

→ decrease scale uncertainty

Work in **narrow width approximation** ($\sim \mathcal{O}(1\%)$ error).

Decays of top quarks and Z-boson include **spin correlations** to NLO.

→ realistic experimental cuts; use spin correlations as discriminating variable

In SM, top-Z coupling is

$$\mathcal{L}_{t\bar{t}Z}^{\text{SM}} = ie\bar{u}(p_t) \left[\gamma^\mu (C_V^{\text{SM}} + \gamma_5 C_A^{\text{SM}}) \right] v(p_{\bar{t}}) Z_\mu$$

$$C_V^{\text{SM}} = \frac{T^3 - 2Q_t \sin^2 \theta_W}{2 \sin \theta_W \cos \theta_W}$$

$$C_A^{\text{SM}} = \frac{-T^3}{2 \sin \theta_W \cos \theta_W}$$

Write **New Physics** in EFT

$$\mathcal{L}_{t\bar{t}Z}^{\text{NP}} = \sum_i \frac{\mathcal{C}_i}{\Lambda^2} O_i + \dots$$

Assuming **dimension-six**, **gauge invariant** operators, Lagrangian is

$$\mathcal{L}_{t\bar{t}Z} = ie\bar{u}(p_t) \left[\gamma^\mu (C_{1,V} + \gamma_5 C_{1,A}) + \frac{i\sigma_{\mu\nu} q_\nu}{M} (C_{2,V} + i\gamma_5 C_{2,A}) \right] v(p_{\bar{t}}) Z_\mu$$

Aguilar-Saavedra, hep-ph/0811.3842

$$\mathcal{L} = ie\bar{u}(p_t) \left[\gamma^\mu (C_{1,V} + \gamma_5 C_{1,A}) + \underbrace{\frac{i\sigma_{\mu\nu}q_\nu}{M} (C_{2,V} + i\gamma_5 C_{2,A})}_{\text{dipole moment}} \right] v(p_{\bar{t}}) Z_\mu$$

Treat $C_{1,V}$, $C_{1,A}$, $C_{2,V}$, $C_{2,A}$ as **anomalous couplings** - independent of kinematics

- Electric and magnetic top dipole moment.
- Zero at tree-level in SM.
- Small loop-induced corrections in SM.
- Non-renormalizable amplitudes.

- Dipole coefficients $C_{2,V}$ and $C_{2,A}$ set to zero.
- Focus on $C_{1,V}$ and $C_{1,A}$.
- Define

$$\Delta C_{1,V} = \frac{C_{1,V}}{C_V^{\text{SM}}} - 1 ; \quad \Delta C_{1,A} = \frac{C_{1,A}}{C_A^{\text{SM}}} - 1.$$

- Calculation performed in TOPAZ (Melnikov, Schulze, ...).
- LO production through $gg \rightarrow t\bar{t}Z$ and $q\bar{q} \rightarrow t\bar{t}Z$.
- Real corrections open qg and $\bar{q}g$ channels.
- Soft and collinear singularities regularized using Catani-Seymour dipoles.
- Virtual corrections to gg and $q\bar{q}$ channels calculated using D -dimensional realization of Ossola-Papadopoulos-Pittau procedure.

Ossola, Papadopoulos, Pittau, hep-ph/0609007; Ellis, Giele, Kunszt, hep-ph/0708.2398;

Giele, Kunszt, Melnikov, hep-ph/0801.2237; Ellis, Giele, Kunszt, Melnikov, hep-ph/0806.3467

Review: Ellis, Kunszt, Melnikov, Zanderighi, hep-ph/1105.4319

- Inclusive cross-section at $\sqrt{s} = 7$ TeV LHC:

$$\sigma_{t\bar{t}Z}^{\text{LO}} = 103.5 \text{ fb};$$

$$\sigma_{t\bar{t}Z}^{\text{NLO}} = 137.0 \text{ fb}$$

(perfect agreement with results of Garzelli, Kardos, Papadopoulos, Trocsanyi)

hep-ph/1111.0610, hep-ph/1111.1444, hep-ph/1208.2665

- In tripletonic decay $t\bar{t}Z \rightarrow (jjbb\nu l^+ l^-)$ at $\sqrt{s} = 13$ LHC TeV:

$$\sigma_{t\bar{t}Z}^{\text{LO}} = 3.80^{+34\%}_{-25\%} \text{ fb};$$

$$\sigma_{t\bar{t}Z}^{\text{NLO}} = 5.32^{+15\%}_{-14\%} \text{ fb}$$

(using $\mu_0 = mt + m_z/2$)

Inclusive cuts:

$$p_{T,j} > 20 \text{ GeV}$$

$$p_{T,l} > 15 \text{ GeV}$$

$$p_{T,\text{miss}} > 20 \text{ GeV}$$

$$|y_l| < 2.5, |y_j| < 2.5$$

$$R_{lj} > 0.4$$


- Scale uncertainty $\pm 28\%$ at LO and $\pm 14\%$ at NLO.
- $k = \sigma^{\text{NLO}} / \sigma^{\text{LO}} \simeq 1.4$.

Compute $d\sigma_{\text{LO}}(\Delta C_{1,V}, \Delta C_{1,A})$ and $d\sigma_{\text{NLO}}(\Delta C_{1,V}, \Delta C_{1,A})$ at $\mu = \mu_0$.
 \Rightarrow large number of computations!


Notice that (at LO or NLO), $\mathcal{A}_{t\bar{t}Z} = A_0 + A_V C_{1,V} + A_A C_{1,A}$.
 Then cross-section

$$d\sigma = s_0 + s_1 C_{1,V} + s_2 C_{1,V}^2 + s_3 C_{1,A} + s_4 C_{1,A}^2 + s_5 C_{1,V} C_{1,A}.$$

- 1 Compute for **six** values of $C_{1,V}$ and $C_{1,A}$.
- 2 Solve for s_i .
- 3 Generate all other values of $d\sigma$.
- 4 Works on **overall cross-sections** and **distributions**.



Dependence of Cross-sections on Top-Z Couplings




- Cross-section changes by approx. 50% in $(\Delta C_{1,V}, \Delta C_{1,A})$ plane.
- Symmetric about $\Delta C_{1,V} = -1$, expected around $\Delta C_{1,A} = -1$.
- Far greater sensitivity to $\Delta C_{1,A}$ than $\Delta C_{1,V}$.
- Cross-sections within scale uncertainty band $\sim 15\%$ **cannot be distinguished** from SM:

e.g. $(\Delta C_{1,V}, \Delta C_{1,A}) = (1.7, -0.3)$

- Use log-likelihood ratio test, with LL ratio derived from Poisson distribution

$$\Lambda(\vec{n}_{\text{obs}}) = \sum_{i=1}^{N_{\text{bins}}} \left[n_{i,\text{obs}} \log\left(\frac{\nu_i^{H_0}}{\nu_i^{H_1}}\right) - \nu_i^{H_0} + \nu_i^{H_1} \right].$$

- $\nu_i^{H_0}$ and $\nu_i^{H_1}$ are *calculated/measured* binned data according to two hypotheses.
- $n_{i,\text{obs}}$ are pseudoexperimental data, generated around one of the hypotheses.
- Generates two distributions for Λ – overlap is a measure of statistical separation of hypotheses.
- Include theoretical uncertainty by uniformly rescaling all bins.



First observation of $t\bar{t}Z$ at the LHC:


ATLAS sees 1 event with 4.7fb^{-1} ,
CMS sees 9 events with 4.9fb^{-1} (bg.
expectation 3.2 events).

\Rightarrow CMS finds

$$\sigma_{t\bar{t}Z} = 0.28^{+0.14}_{-0.11} \text{ (stat.) }^{+0.06}_{-0.03} \text{ (syst.) pb}$$

(Good agreement w.


$$\sigma_{\text{NLO}} = 0.137 \text{ pb.})$$



hep-ex/1303.3239

Use overall cross-section to put **first** direct constraints on top-Z coupling.

- Use Gaussian multiplicative factor for experimental uncertainty (20%).




- SM at $(\Delta C_{1,V}, \Delta C_{1,A}) = (0, 0)$ consistent with measurement [$C_V^{\text{SM}} \simeq 0.24$ and $C_A^{\text{SM}} \simeq -0.60$].
- Rough guide: red excluded at $1\text{-}\sigma$, orange at $2\text{-}\sigma$, yellow at $3\text{-}\sigma$.
- $-11 \lesssim \Delta C_{1,V} \lesssim 10$ and $-4 \lesssim \Delta C_{1,A} \lesssim 2$ at LO (95% C.L.).
- $-8 \lesssim \Delta C_{1,V} \lesssim 7$ and $-3 \lesssim \Delta C_{1,A} \lesssim 1$ at NLO (95% C.L.).
- Much tighter constraints at NLO (reduced scale uncertainty; k -factor).
- But constraints are very loose...

Good start – how can it be improved?


- Better statistics: larger luminosity, higher energy.
- Use **shape information**.

⇒ use $\Delta\phi_{ll}$ distribution at $\sqrt{s} = 13$ TeV for 30, 300, 3000 fb⁻¹.



- Use scale uncertainty 30% at LO and 15% at NLO.
- Obvious improvement with increased luminosity.
- Notable improvement using NLO corrections (reduced scale uncertainty + k -factor).

Focus on 300 fb⁻¹:



- Find $-4.0 < \Delta C_{1,V} < 2.8$ and $-0.36 < \Delta C_{1,A} < 0.54$ at LO.
- At NLO $-3.6 < \Delta C_{1,V} < 1.6$ and $-0.24 < \Delta C_{1,A} < 0.30$.
- $\Rightarrow C_V = 0.24^{+0.39}_{-0.85}$ and $C_A = -0.60^{+0.14}_{-0.18}$ at NLO QCD.

Higher dimensional operators


Higher dimensional operators in EFT \leftrightarrow deviations from SM couplings:

$$C_{1,V} = C_{1,V}^{\text{SM}} + \left(\frac{v^2}{\Lambda^2}\right) \text{Re} \left[C_{\phi q}^{(3,33)} - C_{\phi q}^{(1,33)} - C_{\phi u}^{33} \right],$$

$$C_{1,A} = C_{1,A}^{\text{SM}} + \left(\frac{v^2}{\Lambda^2}\right) \text{Re} \left[C_{\phi q}^{(3,33)} - C_{\phi q}^{(1,33)} + C_{\phi u}^{33} \right],$$

Assuming $SU(2)$ symmetry $\rightarrow C_{\phi q}^{(3,33)} \approx -C_{\phi q}^{(1,33)}$.

Translate constraints on $\Delta C_{1,V}, \Delta C_{1,A}$ into constraints on $C_{\phi q}^{(3,33)}$ and $C_{\phi u}^{33}$:



- $t\bar{t}Z$ production at the LHC calculated to NLO in QCD, including all decays with spin correlation, and using NWA.
- Calculation performed with different values of vector and axial-vector top-Z coupling.
- **Cross-section** compared to that from CMS \rightarrow **first direct detection bounds on top-Z coupling** (very loose).
- Log-likelihood analysis using $\Delta\phi_{ll}$ distribution reveals:
 - \sim factor 2 increase in sensitivity due to decrease in scale uncertainty
 - Couplings giving $\sigma \simeq \sigma_{SM}$ may be distinguished by $\Delta\phi_{ll}$ shape.
 - Constrains $C_V = 0.24^{+0.39}_{-0.85}$ and $C_A = -0.60^{+0.14}_{-0.18}$ at NLO QCD.
- Reduced scale at NLO and $K \simeq 1.5$ boost constraining capability
- Can be related to operators, constrain scale Λ .

Future Work

- Look at constraining coefficients dipole terms $\sim \sigma_{\mu\nu} q^\nu / M$.
- Look at bounds from single top + Z results
- Extend analysis to $t\bar{t} + \gamma$.

Two previous calculations:

Lazopoulos, McElmurry, Melnikov,
Petriello

hep-ph/0804.0610


- No decays
- $\sigma_{\text{LO}} = 0.808 \text{ pb}$, $\sigma_{\text{NLO}} = 1.09 \text{ pb}$,
at $\sqrt{s} = 14 \text{ TeV}$

Garzelli, Kardos, Papadopoulos,
Trocsanyi

hep-ph/1111.0610, hep-ph/1111.1444, hep-ph/1208.2665

- Decays through parton showering,
hadronization effects
- $\sigma_{\text{LO}} = 0.808 \text{ pb}$, $\sigma_{\text{NLO}} = 1.12 \text{ pb}$
at $\sqrt{s} = 14 \text{ TeV}$ (same
parameters)
- $\sigma_{\text{LO}} = 103.5 \text{ fb}$, $\sigma_{\text{NLO}} = 136.9 \text{ fb}$
at $\sqrt{s} = 7 \text{ TeV}$


$\sim 3\%$ tension between results



Same setup as GKPT:


$$\sigma_{\text{LO}}^{\text{RS}} = 103.5\text{fb}; \quad \sigma_{\text{NLO}}^{\text{RS}} = 137.0\text{fb}$$

- **excellent agreement**
at LO and NLO for
both cross-sections
and distributions.



Single top + Z ($tZ + \bar{t}Z$) similar rate to $t\bar{t}Z$

Campbell, Ellis, RR, hep-ph/1302.3856



- Also sensitive to top-Z coupling
- In $tZ \rightarrow b\nu l^+ l^-$ decay, dominant background to tripletonic $t\bar{t}Z$.
- Distinguished by number and behavior of jets.

⇒ defer study of top-Z couplings in single top+Z to later

→ negligible background to $t\bar{t}Z$ signal.

Binned likelihood function with Poisson distribution P_i ,

$$\mathcal{L}(\mathcal{H}|\vec{n}) = \prod_{i=1}^{N_{\text{bins}}} P_i(n_i|\nu_i^{\mathcal{H}}),$$

with n_i events observed and ν_i predicted under hypothesis \mathcal{H} .

Log-likelihoods for predictions under \mathcal{H}_{SM} and \mathcal{H}_{alt} are

$$\log \mathcal{L}(\mathcal{H}_{\text{SM}}, \mathcal{H}_{\text{alt}}|\vec{n}_{\text{obs}}) = \sum_{i=1}^{N_{\text{bins}}} [n_{i,\text{obs}} \log(\nu_i^{\text{SM,alt}}) - \log(n_{i,\text{obs}}!) - \nu_i^{\text{SM,alt}}],$$

and **log-likelihood ratio** is test statistic

$$\begin{aligned} \Lambda(\vec{n}_{\text{obs}}) &= \log \left(\mathcal{L}(\mathcal{H}_{\text{SM}}|\vec{n}_{\text{obs}}) / \mathcal{L}(\mathcal{H}_{\text{alt}}|\vec{n}_{\text{obs}}) \right) \\ &= \sum_{i=1}^{N_{\text{bins}}} \left[n_{i,\text{obs}} \log \left(\frac{\nu_i^{\text{SM}}}{\nu_i^{\text{alt}}} \right) - \nu_i^{\text{SM}} + \nu_i^{\text{alt}} \right], \end{aligned}$$

$$\Lambda(\vec{n}_{\text{obs}}) = \sum_{i=1}^{N_{\text{bins}}} \left[n_{i,\text{obs}} \log\left(\frac{\nu_i^{\text{SM}}}{\nu_i^{\text{alt}}}\right) - \nu_i^{\text{SM}} + \nu_i^{\text{alt}} \right],$$


Use **pseudoexperimental data** for \vec{n}_{obs} .

- Generated using bin-by-bin Poisson distribution around $\vec{\nu}^{\text{SM}}$
- Repeat many times \rightarrow **distribution** $P(\Lambda|\mathcal{H}_{\text{SM}})$.
- Now generate $P(\Lambda|\mathcal{H}_{\text{alt}})$ by using $\vec{\nu}^{\text{alt}}$ to generate \vec{n}_{obs} .
- Overlap of $P(\Lambda|\mathcal{H}_{\text{SM}})$ and $P(\Lambda|\mathcal{H}_{\text{alt}})$ gives statistical separation of hypotheses

Type-I error (falsely reject \mathcal{H}_{alt}) and type-II error (falsely reject \mathcal{H}_{SM}) given by

$$\alpha = \int_{\hat{\Lambda}}^{\infty} d\Lambda P(\Lambda|\mathcal{H}_{\text{SM}})$$

$$\beta = \int_{-\infty}^{\hat{\Lambda}} d\Lambda P(\Lambda|\mathcal{H}_{\text{alt}}).$$



De Rújula et. al., hep-ph/1001.5300

We choose $\alpha = \beta$ – equal chance of incorrectly rejecting each hypothesis in favor of the other.

Can convert to sigma-level through

$$\sigma = \sqrt{2} \operatorname{erf}^{-1}(1 - \alpha),$$

Three operators involved in top-Z coupling:

$$\mathcal{O}_{\phi q}^{(1)} = i(\phi^\dagger D_\mu \phi)(\bar{q}\gamma^\mu q)$$

$$\mathcal{O}_{\phi q}^{(3)} = i(\phi^\dagger \tau^I D_\mu \phi)(\bar{q}\gamma^\mu q)$$

$$\mathcal{O}_{\phi t} = i(\phi^\dagger D_\mu \phi)(\bar{t}_R \gamma^\mu t_R)$$




and

$$\delta C_L = \text{Re}(C_{\phi q}^{(3)} - C_{\phi q}^{(1)}) \frac{v^2}{\Lambda^2}$$

$$\delta C_R = -\text{Re} C_{\phi t} \frac{v^2}{\Lambda^2}$$


Aguilar-Saavedra, hep-ph/0811.3842; Berger, Cao, Low, hep-ph/0907.2191

- $C_{\phi q}^{(3)} + C_{\phi q}^{(1)}$ tightly constrained by $Z \rightarrow b\bar{b}$ (assuming $SU(2)_L \times U(1)_Y$ symmetry).
- $t \rightarrow Wb$ depends on $C_{\phi q}^{(3)} - C_{\phi q}^{(1)}$ and $|V_{tb}|$
- Accurate measurement of top-Z coupling in $t\bar{t}Z \rightarrow$ get $|V_{tb}|$



Study by Denner, Dittmaier, Kallweit, Pozzorini, Schulze, for SM NLOWG, hep-ph/1203.6803


- Shape changes when including decay
- No difference using NWA vs. full calculation for $m_{lb} \lesssim 2m_t$.
- Shape changes $m_{lb} \simeq 2m_t$.



Baur, Juste, Orr,

Rainwater:

hep-ph/0412021



Baur, Juste, Orr,
 Rainwater;
 hep-ph/0412021