

RCNP E398 experiment $C,O(p,p')$ to measure γ ray branching ratio ($E > 5 \text{ MeV}$) from the giant resonances of carbon and oxygen in relation to the γ ray production in $C,O(\nu, \nu')$.

Friday, 12 September 2014 18:00 (1h 30m)

We plan to measure the branching ratios of γ -ray emission ($E_\gamma > 5 \text{ MeV}$) from giant resonance of ^{16}O and ^{12}C , as the functions of excitation energy (E_x).

This measurement will provide the fundamental and important information not only for the γ -ray production from primary neutral-current neutrino-oxygen (-carbon) interactions but also for that from the secondary hadronic (neutron-oxygen and -carbon) interactions.

The understanding of the γ -ray production will introduce a new neutrino detection method to Supernova neutrino physics and Neutrino oscillation physics.

In the second stage, we would like to perform $O,C(\text{He},t)$ ($T=1$) experiment at 0 degrees to continue the systematic study of spin-isospin response through the measurement of the γ -ray production with oxygen and carbon nuclei.

Ref.

[1] T.Mori, M.Sakuda, A.Tamii, H.Toki, M.Nakahata, and K.Ueno, Study of γ -ray production

from neutral-current neutrino-Oxygen interaction and the detection of the neutrino

from Supernova explosion, AIP Conf. Proc.1269, 418-420, 2010.

[2] A.Ankowski, O.Benhar, T.Mori, R.Yamaguchi, and M.Sakuda,

Analysis of γ -ray production in neutral-current neutrino-oxygen quasi-elastic interactions above 200 MeV,

Phys.Rev.Lett.108,052505(2012).

Primary author: Mr OU, Iwa (Okayama University)

Co-authors: Prof. SAKUDA, Makoto (Okayama University); Mr MORI, Takaaki (Okayama University); Dr YANO, Takatomi (Okayama University)

Presenter: Mr OU, Iwa (Okayama University)

Session Classification: Happy hour with posters

Track Classification: Happy hour with posters