



Contribution ID: 180

Type: Poster Presentation

First direct determination of the superallowed β -decay Q-EC-value for O-14 via Penning trap mass spectrometry at the LEBIT facility

Monday, 11 May 2015 16:30 (1 minute)

The conserved vector current (CVC) hypothesis asserts that the vector part of the weak interaction is independent of the nuclear interaction. This means that the vector coupling constant G_V is truly a constant and does not require renormalization. This constant, when combined with the purely leptonic muon decay constant G_F determines the up-down matrix element V_{ud} of the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix. Measurements of the ft values of super-allowed $0^+ \rightarrow 0^+$ transitions allow G_V to be determined. A precise determination of V_{ud} has been, and continues to be, an important component of the test of the unitarity of the CKM quark-mixing matrix and the search for physics beyond the standard model. To determine an ft value, the decay half-life, the branching ratio for the $0^+ \rightarrow 0^+$ transition, and the QEC values need to be known.

Low-Z, superallowed β -emitters like O-14 are particularly significant for setting limits on the existence of scalar currents. While the CVC hypothesis states that F_t should be the same for all superallowed $0^+ \rightarrow 0^+$ β -decays, if there is a scalar interaction, an additional term approximately inversely proportional to QEC would be present in F_t . As QEC-values are smaller for lower-Z isotopes, these isotopes would be most sensitive to the presence of a scalar current, showing the largest deviation in F_t . To date, 14 F_t values are used to calculate the world average [1]. Of these 14 decays, only O-14 has not been measured in a Penning trap, despite multiple attempts at other facilities. At LEBIT [2] we have performed the first direct measurement of the ground state β -decay QEC value. This measurement provides an order of magnitude improvement in precision, and it no longer makes a significant contribution to the uncertainty of its associated F_t value. Together with future reductions in the uncertainties of the other major contributors, particularly the half life and branching ratio, more stringent limits will be placed on the existence of scalar currents in the electroweak interaction.

[1] J. C. Hardy and I. S. Towner, Phys. Rev. C 91, 025501 (2015).

[2] R. J. Ringle, S. Schwarz, and G. Bollen, Int. J. Mass Spectrom. 349-350, 87 (2013).

Primary author: Ms SANDLER, Rachel (National Superconducting Cyclotron Laboratory/ Michigan State University)

Co-authors: Mr VALVERDE, Adrian (Michigan State University / National Superconducting Cyclotron Laboratory); Dr VILLARI, Antonio C.C. (Facility for Rare Isotope Beams - MSU); Dr SUMITHRARACHCHI, Chandana (National Superconducting Cyclotron Laboratory); Mr IZZO, Christopher (National Superconducting Cyclotron Laboratory/ Michigan State University); Mr MORRISSEY, David (MSU/NSCL); BOLLEN, Georg (Michigan State University); Dr GULYUZ, Kerim (NSCL); Ms COOPER, Kortney (MSU/NSCL); Dr EIBACH, Martin (National Superconducting Cyclotron Laboratory); Dr REDSHAW, Matthew (Central Michigan University); Prof. BRODEUR, Maxime (University of Notre Dame); BRYCE, Richard Adam (Central Michigan University); Dr RINGLE, Ryan (NSCL); Dr SCHWARZ, Stefan (NSCL)

Presenter: Ms SANDLER, Rachel (National Superconducting Cyclotron Laboratory/ Michigan State University)

Session Classification: Poster Session A