

# Resummation of double-differential cross sections

based on M. Procura, W. J. Waalewijn and L. Z., JHEP 1502 (2015) 117



Lisa Zeune

University of Amsterdam

SCET 2015

Santa Fe, 26th March 2015



UNIVERSITEIT VAN AMSTERDAM



# Motivation

Why shall we study multi-differential cross sections?

- LHC analyses often involve several measurements/cuts → See talks by Andrew Larkoski (SCET 2014), Ian Mout and Piotr Pietrulewicz (both today)
- Correlations between different observables encoded in multi-differential cross sections

## Example:

Z + 0 jet: Global jet veto using beam thrust and measurement of the transverse momentum of the Z boson

- If the measurements lead to widely separated energy scales → resummation required
- So far: resummed calculation (mostly) restricted to single variables  
Resummation of multi-differential cross sections have been studied — but there the measurements concerned different regions of phase-space (e.g different jets).  
E.g. Ellis, Vermilion, Walsh, Hornig, Lee, '10; Stewart, Tackmann, Waalewijn, '10; Kelley, Schwartz, Schabinger, Zhu, '11; for a discussion of NGLs, see talk by Duff Neill

# Motivation

- Another important reason to study the resummation of multi/double - differential cross sections: **Jet substructure**
  - ↳ One goal: Discriminate QCD jets from heavy boosted particles (W, Z, H, t)
- Most powerful discrimination observables are ratios of infrared and collinear (IRC) safe observables

## Examples:

Ratios of N-subjettiness, energy correlation functions or planar flow /  
Ratio of two angularities

Resummation of this ratio observable was studied in JHEP 1409 (2014) 046 (Larkoski, Moult, Neill), which inspired our work

- These observables are not IRC safe (cannot be computed order-by-order in  $\alpha_S$ ), but can be calculated in a well-defined way by marginalising over the **resummed** double differential cross section. [Larkoski, Thaler, '13](#)

# Outline

In this talk I will present an extension of SCET which enables the resummation of a class of double-differential measurements

- **Application 1:  $Z + 0$  jet production**

- Introduction to SCET+
- Factorization formula

- **Application 2: Measurement of two angularities on a jet**

- NLL cross section
- Comparison to JHEP 1409 (2014) 046 (Larkoski, Moult, Neill)

# Outline

In this talk I will present an extension of SCET which enables the resummation of a class of double-differential measurements

- **Application 1:  $Z + 0$  jet production**
  - Introduction to SCET+
  - Factorization formula

# Introduction to SCET+

- Consider  $Z + 0$  jet production:

Transverse momentum of  $Z$  measured

and global jet veto imposed using beam thrust  $\mathcal{T}$

$$\begin{aligned}\mathcal{T} &= \sum_i p_{iT} e^{-|\eta_i|} \\ &= \sum_i \min\{p_i^+, p_i^-\}\end{aligned}$$

Stewart, Tackmann,  
Waalewijn, '09

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

→ **SCET I:**

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$


→ **SCET+:**

$$p_T \sim Q^{1-r} \mathcal{T}^r$$

with  $1/2 < r < 1$

→ **SCET II:**

$$p_T \sim \mathcal{T}$$



# Introduction to SCET+

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

## SCET I:

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$

## SCET+:

$$p_T \sim Q^{1-r} \mathcal{T}^r$$

with  $1/2 < r < 1$

## SCET II:

$$p_T \sim \mathcal{T}$$


Mode	Scaling $(-, +, \perp)$	Measurement
$n$ -collinear	$Q(1, \lambda^2, \lambda)$	$p_T \sim Q\lambda$
soft	$Q(\lambda^2, \lambda^2, \lambda^2)$	$\mathcal{T} \sim Q\lambda^2$

Fully-unintegrated (FU) beam functions:

$B_q(t, x, \vec{k}_\perp)$  → momentum fraction  
 →  $-t = k^- k^+$ : transverse virtuality

Soft function:

$$S(k^+)$$



Interactions removed by BPS field redefinition

Bauer, Pirjol, Stewart, '02



# Introduction to SCET+

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

**SCET I:**

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$

**SCET+:**


$$p_T \sim Q^{1-r} \mathcal{T}^r$$

with  $1/2 < r < 1$

**SCET II:**

$$p_T \sim \mathcal{T}$$

Mode	Scaling ( $-$ , $+$ , $\perp$ )	Measurement
$n$ -collinear	$Q(1, \lambda^2, \lambda)$	$p_T \sim Q\lambda$
soft	$Q(\lambda, \lambda, \lambda)$	$\mathcal{T} \sim Q\lambda$



**TMD beam functions:**

$$B_q(x, \vec{k}_\perp)$$

**FU soft function:**

$$S(k^+, \vec{k}_\perp)$$



# Introduction to SCET+

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

**SCET I:**

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$

**SCET+:**

$$p_T \sim Q^{1-r} \mathcal{T}^r$$


with  $1/2 < r < 1$

**SCET II:**

$$p_T \sim \mathcal{T}$$

Mode	Scaling ( $-$ , $+$ , $\perp$ )	Measurement
$n$ -collinear	$Q(1, \lambda^{2r}, \lambda^r)$	$p_T$
$n$ -collinear-soft	$Q(\lambda^{2r-1}, \lambda, \lambda^r)$	$\mathcal{T}$
soft	$Q(\lambda, \lambda, \lambda)$	$\mathcal{T}$

with  $1/2 < r < 1$ ,  
 $\lambda \sim \mathcal{T}/Q \sim (p_T/Q)^{1/r}$



**TMD beam functions:**

$$B_q(x, \vec{k}_\perp)$$

**Soft function:**

$$S(k^+)$$

**Collinear-soft function:**

$$\mathcal{S}(k^+, \vec{k}_\perp)$$

Modes need to be well separated for power corrections to be small

# Introduction to SCET+

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

**SCET I:**

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$

**SCET+:**

$$p_T \sim Q^{1-r} \mathcal{T}^r$$


with  $1/2 < r < 1$

**SCET II:**

$$p_T \sim \mathcal{T}$$

Mode	Scaling ( $-$ , $+$ , $\perp$ )	Measurement
$n$ -collinear	$(Q, p_T^2/Q, p_T)$	$p_T$
$n$ -collinear-soft	$(p_T^2/\mathcal{T}, \mathcal{T}, p_T)$	$\mathcal{T}$
soft	$(\mathcal{T}, \mathcal{T}, \mathcal{T})$	$\mathcal{T}$

with  $1/2 < r < 1$ ,  
 $\lambda \sim \mathcal{T}/Q \sim (p_T/Q)^{1/r}$



**TMD beam functions:**

$$B_q(x, \vec{k}_\perp)$$

**Soft function:**

$$S(k^+)$$

**Collinear-soft function:**

$$\mathcal{S}(k^+, \vec{k}_\perp)$$

Modes need to be well separated for power corrections to be small

# Introduction to SCET+

Hierarchy between  $\mathcal{T}$  and  $p_T$  determines the appropriate SCET version:

SCET I:

$$p_T \sim Q^{1/2} \mathcal{T}^{1/2}$$

SCET+:

$$p_T \sim Q^{1-r} \mathcal{T}^r$$


with  $1/2 < r < 1$

SCET II:

$$p_T \sim \mathcal{T}$$

- Collinear-soft modes were introduced first in a different context in Phys.Rev.D85 (2012) 074006 (Bauer, Tackmann, Walsh, Zuberi) and has led us to adopt their name SCET+

→ Difference: In their case collinear and collinear-soft modes are separated in virtuality (SCET I like) while in our case collinear and collinear-soft modes are separated in rapidity (SCET II like)



# Effective theory framework

## I. Matching the QCD quark current onto SCET+

$$\bar{\Psi} \Gamma \Psi = C(Q^2, \mu) \bar{\xi}_{\bar{n}} W_{\bar{n}}^\dagger S_{\bar{n}}^\dagger X_{\bar{n}}^\dagger V_{\bar{n}} \Gamma V_n^\dagger X_n S_n W_n^\dagger \xi_n$$

QCD quark fields  $\uparrow$  Dirac structure  $\uparrow$  matching coefficient  $\uparrow$  collinear antiquark moving in  $\bar{n}$ -direction  $\uparrow$  collinear quark moving in  $n$ -direction

### Wilson lines

$W_n^\dagger$ :  $n$ -collinear gluons emitted from  $\bar{\Psi}$  ( $\bar{n}$ -collinear)

$V_n^\dagger$ :  $n$ -collinear-soft gluons emitted from  $\bar{\Psi}$  ( $\bar{n}$ -collinear)

$S_n$ : soft gluons emitted from  $\Psi$  ( $n$ -collinear)

$X_n$ :  $n$ -collinear-soft gluons emitted from  $\Psi$  ( $n$ -collinear)

The ordering of the Wilson lines is fixed by gauge invariance of SCET+

# Effective theory framework

## II. BPS field redefinition

- At this point the soft fields still interact with the collinear-soft fields
- Performing an analog to the BPS field redefinition: Bauer, Pirjol, Stewart, '02

$$V_n \rightarrow S_n V_n S_n^\dagger,$$

$$X_n \rightarrow S_n X_n S_n^\dagger,$$


$$V_{\bar{n}} \rightarrow S_{\bar{n}} V_{\bar{n}} S_{\bar{n}}^\dagger,$$

$$X_{\bar{n}} \rightarrow S_{\bar{n}} X_{\bar{n}} S_{\bar{n}}^\dagger$$

- Finally:

$$\bar{\Psi} \Gamma \Psi = C(Q^2, \mu) \bar{\xi}_{\bar{n}} W_{\bar{n}} X_{\bar{n}}^\dagger V_{\bar{n}} S_{\bar{n}}^\dagger \Gamma S_n V_n^\dagger X_n W_n^\dagger \xi_n$$

- No interaction between various modes anymore  
→ Derive factorisation theorems



# Factorisation theorems: SCET I

Stewart, Tackmann, Waalewijn, '09;  
Jain, Procura, Waalewijn, '11

$$\frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} = \sum_q \hat{\sigma}_q^0 H(Q^2) \int dt_1 dt_2 \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} \int dk^+ S(k^+) \\ \times \left[ B_q(t_1, x_1, \vec{k}_{1\perp}) B_{\bar{q}}(t_2, x_2, \vec{k}_{2\perp}) + (q \leftrightarrow \bar{q}) \right] \\ \times \delta\left(\mathcal{T} - \frac{e^{-Y} t_1 + e^Y t_2}{Q} - k^+\right) \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp}|^2)$$


- Ingredients:

## FU beam function

$\mathbf{P}$  operator returns momentum  
of intermediate state

$$\chi_n = W_n^\dagger \xi_n$$

$$B_q(t, x, \vec{k}_\perp) = \left\langle p_n(p^-) \left| \bar{\chi}_n(0) \frac{\not{n}}{2} \left[ \delta(k^- - p^- + \mathbf{P}^-) \right. \right. \right. \\ \left. \left. \left. \delta(t - k^- \mathbf{P}^+) \delta^2(\vec{k}_\perp - \vec{\mathbf{P}}_\perp) \chi_n(0) \right] \right| p_n(p^-) \right\rangle$$



$\hat{\sigma}_q^0$ : Born cross section

$H(Q^2, \mu) = |C(Q^2, \mu)|^2$ : Hard function

$-t_i = k_i^- k_i^+$  ( $i = 1, 2$ ): Transverse virtuality

$x_i = Q/E_{\text{cm}} e^{\pm Y}$  ( $i = 1, 2$ ): Momentum fraction,  
 $Y = \text{rapidity}$

# Factorisation theorems: SCET I

Stewart, Tackmann, Waalewijn, '09;  
Jain, Procura, Waalewijn, '11


$$\begin{aligned} \frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} &= \sum_q \hat{\sigma}_q^0 H(Q^2) \int dt_1 dt_2 \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} \int dk^+ S(k^+) \\ &\times \left[ B_q(t_1, x_1, \vec{k}_{1\perp}) B_{\bar{q}}(t_2, x_2, \vec{k}_{2\perp}) + (q \leftrightarrow \bar{q}) \right] \\ &\times \delta\left(\mathcal{T} - \frac{e^{-Y} t_1 + e^Y t_2}{Q} - k^+\right) \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp}|^2) \end{aligned}$$

- Ingredients:

## Soft function

$$S(k^+) = \frac{1}{N_c} \langle 0 | \text{Tr} \left[ \overline{\mathbf{T}}(S_n^\dagger(0) S_{\bar{n}}(0)) \delta(k^+ - \mathbf{P}_1^+ - \mathbf{P}_2^-) \mathbf{T}(S_{\bar{n}}^\dagger(0) S_n(0)) \right] | 0 \rangle$$

Time ordering



$\mathbf{P}_1$  operator returns momentum of soft radiation in hemisphere 1 ( $p^+ < p^-$ )

$\hat{\sigma}_q^0$  : Born cross section

$H(Q^2, \mu) = |C(Q^2, \mu)|^2$  : Hard function

$-t_i = k_i^- k_i^+$  ( $i = 1, 2$ ) : Transverse virtuality

$x_i = Q/E_{\text{cm}} e^{\pm Y}$  ( $i = 1, 2$ ) : Momentum fraction,  
 $Y = \text{rapidity}$



# Factorisation theorems: SCET II

$$\frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} = \sum_q \hat{\sigma}_q^0 H(Q^2) \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} d^2\vec{k}_\perp \int dk^+ \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp} + \vec{k}_\perp|^2) \\ \times \delta(\mathcal{T} - k^+) \left[ B_q(x_1, \vec{k}_{1\perp}) B_{\bar{q}}(x_2, \vec{k}_{2\perp}) + (q \leftrightarrow \bar{q}) \right] S(k^+, \vec{k}_\perp)$$

Extension of: Chiu, Jain, Neill, Rothstein, '12  
See also: Becher, Neubert, '10

- Ingredients:

## TMD beam function

$$B_q(x, \vec{k}_\perp) = \left\langle p_n(p^-) \left| \bar{\chi}_n(0) \frac{\not{n}}{2} \left[ \delta(k^- - p^- + \mathbf{P}^-) \delta^2(\vec{k}_\perp - \vec{\mathbf{P}}_\perp) \chi_n(0) \right] \right| p_n(p^-) \right\rangle$$

## FU soft function

$$S(k^+, \vec{k}_\perp) = \frac{1}{N_c} \langle 0 | \text{Tr} \left[ \mathbf{T} (S_n^\dagger(0) S_{\bar{n}}(0)) \delta(k^+ - \mathbf{P}_1^+ - \mathbf{P}_2^-) \right. \\ \left. \delta^2(\vec{k}_\perp - \vec{\mathbf{P}}_\perp) \mathbf{T} (S_{\bar{n}}^\dagger(0) S_n(0)) \right] | 0 \rangle$$

# Factorisation theorems: SCET+

$$\begin{aligned} \frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} &= \sum_q \hat{\sigma}_q^0 H(Q^2) \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} d^2\vec{k}_{1\perp}^{\text{cs}} d^2\vec{k}_{2\perp}^{\text{cs}} \int dk_1^+ dk_2^+ dk^+ \\ &\times S(k^+) B_q(x_1, \vec{k}_{1\perp}) B_{\bar{q}}(x_2, \vec{k}_{2\perp}) \\ &\times \mathcal{J}(k_1^+, \vec{k}_{1\perp}^{\text{cs}}) \mathcal{J}(k_2^+, \vec{k}_{2\perp}^{\text{cs}}) \delta(\mathcal{T} - k_1^+ - k_2^+ - k^+) \\ &\times \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp} + \vec{k}_{1\perp}^{\text{cs}} + \vec{k}_{2\perp}^{\text{cs}}|^2) + (q \leftrightarrow \bar{q}) \end{aligned}$$

- Ingredients:

Soft function  $\rightarrow$  SCET I

TMD beam function  $\rightarrow$  SCET II

- In SCET+ we have a TMD beam function without a TMD soft function

We cannot combine them as was done in Becher, Neubert, '11; Echevarria, Idilbi, Scimemi, '12

# Factorisation theorems: SCET+

$$\begin{aligned} \frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} &= \sum_q \hat{\sigma}_q^0 H(Q^2) \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} d^2\vec{k}_{1\perp}^{\text{cs}} d^2\vec{k}_{2\perp}^{\text{cs}} \int dk_1^+ dk_2^+ dk^+ \\ &\times S(k^+) B_q(x_1, \vec{k}_{1\perp}) B_{\bar{q}}(x_2, \vec{k}_{2\perp}) \\ &\times \mathcal{S}(k_1^+, \vec{k}_{1\perp}^{\text{cs}}) \mathcal{S}(k_2^+, \vec{k}_{2\perp}^{\text{cs}}) \delta(\mathcal{T} - k_1^+ - k_2^+ - k^+) \\ &\times \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp} + \vec{k}_{1\perp}^{\text{cs}} + \vec{k}_{2\perp}^{\text{cs}}|^2) + (q \leftrightarrow \bar{q}) \end{aligned}$$




- Ingredients:

Collinear-soft functions (separately for  $n$  and  $\bar{n}$  directions)

$$\begin{aligned} \mathcal{S}(k^+, \vec{k}_\perp) &= \frac{1}{N_c} \langle 0 | \text{Tr} [\overline{\mathbf{T}}(X_n^\dagger(0) V_n(0)) \delta(k^+ - \mathbf{P}^+) \delta^2(\vec{k}_\perp - \vec{\mathbf{P}}_\perp) \mathbf{T}(V_n^\dagger(0) X_n(0))] | 0 \rangle \\ &= \frac{1}{N_c} \langle 0 | \text{Tr} [\overline{\mathbf{T}}(V_{\bar{n}}^\dagger(0) X_{\bar{n}}(0)) \delta(k^+ - \mathbf{P}^-) \delta^2(\vec{k}_\perp - \vec{\mathbf{P}}_\perp) \mathbf{T}(X_{\bar{n}}^\dagger(0) V_{\bar{n}}(0))] | 0 \rangle \end{aligned}$$

- FU soft function and collinear-soft function look quite similar  
Difference: Collinear-soft radiation goes only into one hemisphere  
 → Different treatment of the two hemispheres

# Summary factorization theorems



$$d\sigma = H(Q^2) \times B_q(t_1, \vec{k}_{1\perp}) B_{\bar{q}}(t_2, \vec{k}_{2\perp}) \times S(k^+)$$

$$d\sigma = H(Q^2) \times B_q(\vec{k}_{1\perp}) B_{\bar{q}}(\vec{k}_{2\perp}) \times \mathcal{S}(k_1^+, \vec{k}_{1\perp}^{\text{cs}}) \mathcal{S}(k_2^+, \vec{k}_{2\perp}^{\text{cs}}) \times S(k^+)$$

$$d\sigma = H(Q^2) \times B_q(\vec{k}_{1\perp}) B_{\bar{q}}(\vec{k}_{2\perp}) \times S(k^+, \vec{k}_{\perp})$$

# Matching of the effective theories

- The **SCET I**, **SCET+** and **SCET II** factorization theorems can be matched achieving a continuous cross section description

**SCET I** ← **SCET+**

$$\mathcal{I}_{ij}(t, x, \vec{k}_\perp) = \int d^2\vec{k}'_\perp \mathcal{I}_{ij}(x, \vec{k}'_\perp) \mathcal{S}(t/p^-, \vec{k}_\perp - \vec{k}'_\perp) + \text{power corrections}$$

beam function matching coefficients\*

$$S(k^+, \vec{k}_\perp) = \int d^2\vec{k}'_\perp \int dk'^+ dk''^+ S(k^+ - k'^+ - k''^+) \mathcal{S}(k'^+, \vec{k}'_\perp) \mathcal{S}(k''^+, \vec{k}_\perp - \vec{k}'_\perp) + \text{power corrections}$$

**SCET II** ← **SCET+**

This holds for common scales:  $\mu = \mu_B = \mu_{\mathcal{S}} = \mu_S$  and  $\nu = \nu_B = \nu_{\mathcal{S}} = \nu_S$

- This follows from:
  - Switching off resummation, SCET I and SCET II produce fixed order cross section up to power corrections
  - SCET+ regime can be obtained by a further expansion of SCET I or SCET II

$$*B_q(x, \vec{k}_\perp, \mu, \nu) = \sum_j \int_x^1 \frac{dx'}{x'} \mathcal{I}_{qj}\left(\frac{x}{x'}, \vec{k}_\perp, \mu, \nu\right) f_j(x', \mu) \left[1 + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}^2}{\vec{k}_\perp^2}\right)\right]$$

# NNLL resummation and consistency checks

- All ingredients entering the factorisation calculated to the accuracy needed for NNLL resummation
  - ↳ **New pieces:** FU soft function and collinear soft function, both calculated at one-loop
  - ↳ No more details here → see paper

## Checks of our framework

- Cancellation of anomalous dimensions between the various ingredients shown
- NLO cross section:
  - Full NLO cross section (differential in  $Q^2$ ,  $Y$ ,  $p_T$  and  $\mathcal{T}$ ) calculated
  - Expanded in the SCET I, SCET+ and SCET II regions of phase space
  - Agreement with the predictions from factorization theorems shown

# Outline

In this talk I will present an extension of SCET which enables the resummation of a class of double-differential measurements

- **Application 2: Measurement of two angularities on a jet**
  - NLL cross section
  - Comparison to JHEP 1409 (2014) 046 (Larkoski, Moult, Neill)



# Measuring two angularities on one jet

- Definition of angularities:

Almeida, Lee, Perez, Sterman, Sung, Virzi, '09;  
Ellis, Vermilion, Walsh, Hornig, Lee, '10;  
Berger, Kucs, Sterman, '03


$$e_\alpha = \frac{1}{E_J} \sum_{i \in J} E_i \frac{\sin \theta_i \tan^{\alpha-1} \frac{\theta_i}{2}}{\sin R \tan^{\alpha-1} \frac{R}{2}} \approx \frac{1}{E_J} \sum_{i \in J} E_i \left( \frac{\theta_i}{R} \right)^\alpha$$

Jet energy

Approximation valid for  $R \ll 1$


$\alpha > 0$

for IRC safety



recoil free  
jet axis

Larkoski, Neill, Thaler, '14;  
Bertolini, Chan, Thaler, '13



- Phase space for the measurement of two angularities  $e_\alpha$  and  $e_\beta$  between:

**Boundary B1:**  $e_\alpha = e_\beta$

(from jet radius requirement)


$\alpha > \beta$

**Boundary B2:**  $e_\alpha^\beta = e_\beta^\alpha$

(from energy conservation)

# Larkoski, Moul, Neill: NLL conjecture

- Two boundary theories for the measurement of two angularities on a single jet were identified
- Factorization of the double differential cross section proven at the phase space boundaries
- Interpolating function across the bulk region derived






Larkoski, Moul,  
Neill, '14

→ requiring cumulative cross section to be continuous and have a continuous derivative at the boundaries

# Double angularities in SCET+

- SCET+ can be used to describe bulk region

Mode	Scaling $(-, +, \perp)$	Measurement
$n$ -collinear	$Q(1, \lambda^{2r/\beta}, \lambda^{r/\beta})$	 $e_\beta$
$n$ -collinear-soft	$Q\left(\lambda^{\frac{\alpha r - \beta}{\alpha - \beta}}, \lambda^{\frac{(\alpha - 2)r - (\beta - 2)}{\alpha - \beta}}, \lambda^{\frac{(\alpha - 1)r - (\beta - 1)}{\alpha - \beta}}\right)$	
soft	$Q(\lambda, \lambda, \lambda)$	 $e_\alpha$

$\beta/\alpha < r < 1$   
and  $\lambda \sim e_\alpha \sim e_\beta^{1/r}$

- Factorization formula (valid to NLL)

$$\frac{d^2\sigma_i}{de_\alpha de_\beta} = \hat{\sigma}_i^{(0)} H_i(Q^2) \int de_\beta^c Q^\beta de_\alpha^{cs} Q de_\beta^{cs} Q^\beta de_\alpha^s Q$$

$\uparrow$   
 $i = q$  (quarks)  
 $i = g$  (gluons)

$J_i(e_\beta^c Q^\beta) \mathcal{S}_i(e_\alpha^{cs} Q, e_\beta^{cs} Q^\beta) S_i(e_\alpha^s Q)$

$$\times \delta(e_\alpha - e_\alpha^{cs} - e_\alpha^c) \delta(e_\beta - e_\beta^c - e_\beta^{cs})$$

(avoiding  $\alpha = 1$  or  $\beta = 1$ )

# RG equations

## Hard function

$$\mu \frac{d}{d\mu} H(Q^2, \mu) = \gamma_H(Q^2, \mu) H(Q^2, \mu),$$

non-cusp piece  
 $\gamma_X^i(\alpha_s) = \sum_n \gamma_{X,n}^i \left(\frac{\alpha_s}{4\pi}\right)^{n+1}$

$$\gamma_H(Q^2, \mu) = \Gamma_{\text{cusp}}(\alpha_s) \ln \frac{Q^2}{\mu^2} + \gamma_H(\alpha_s)$$

## Jet function

cusp piece

$$\mu \frac{d}{d\mu} J(e_\beta Q^\beta, \mu) = \int_0^{e_\beta} de'_\beta Q^\beta \gamma_J(e_\beta Q^\beta - e'_\beta Q^\beta, \mu) J(e'_\beta Q^\beta, \mu),$$
$$\gamma_J(e_\beta Q^\beta, \mu) = -\frac{2}{\beta-1} \Gamma_{\text{cusp}}(\alpha_s) \frac{1}{\mu^\beta} \mathcal{L}_0\left(\frac{e_\beta Q^\beta}{\mu^\beta}\right) + \gamma_J(\alpha_s) \delta(e_\beta Q^\beta)$$

## Soft function

$$\mu \frac{d}{d\mu} S(e_\alpha Q, \mu) = \int_0^{e_\alpha} de'_\alpha Q \gamma_S(e_\alpha Q - e'_\alpha Q, \mu) S(e'_\alpha Q, \mu),$$
$$\gamma_S(e_\alpha Q, \mu) = \frac{2}{\alpha-1} \Gamma_{\text{cusp}}(\alpha_s) \frac{1}{\mu} \mathcal{L}_0\left(\frac{e_\alpha Q}{\mu}\right) + \gamma_S(\alpha_s) \delta(e_\alpha Q)$$

## Collinear-soft function constrained by consistency

# NLL resummation

- Tree level expressions

$$H(Q^2, \mu) = 1,$$

$$J(e_\beta Q^\beta, \mu) = \delta(e_\beta Q^\beta)$$

$$\mathcal{S}(e_\alpha Q, e_\beta Q^\beta) = \delta(e_\alpha Q) \delta(e_\beta Q^\beta),$$

$$S(e_\alpha Q, \mu) = \delta(e_\alpha Q)$$

- Evolve all to the collinear-soft scale  $\mu_{\mathcal{S}}$ :

## Double cumulative distribution

$$\begin{aligned} \Sigma(e_\alpha, e_\beta) &= \int_0^{e_\alpha} de'_\alpha \int_0^{e_\beta} de'_\beta \frac{\partial^2 \sigma}{\partial e'_\alpha \partial e'_\beta} \\ &= \hat{\sigma}^{(0)} \frac{e^{K_H + K_J + K_S - \gamma_E \eta_J - \gamma_E \eta_S}}{\Gamma(1 + \eta_J) \Gamma(1 + \eta_S)} \left( \frac{Q}{\mu_H} \right)^{2\eta_H} \left( \frac{e_\beta^{1/\beta} Q}{\mu_J} \right)^{\beta \eta_J} \left( \frac{e_\alpha Q}{\mu_S} \right)^{\eta_S} \end{aligned}$$

Hard scale

Jet scale

Soft scale

- Evolution kernels:  $K_X(\mu_X, \mu_{\mathcal{S}})$  and  $\eta_X(\mu_X, \mu_{\mathcal{S}})$ ,  $X = H, J, S$

# Comparison to Larkoski, Moulton, Neill

- Their NLL conjecture:

$$\Sigma(e_\alpha, e_\beta)^{\text{conjecture}} = \frac{e^{-\gamma_E \tilde{R}(e_\alpha, e_\beta)}}{\Gamma(1 + \tilde{R}(e_\alpha, e_\beta))} e^{-R(e_\alpha, e_\beta) - \gamma_i T(e_\alpha, e_\beta)}$$

- This mostly agrees with our result with

$$R(e_\alpha, e_\beta) + \gamma T(e_\alpha, e_\beta) \stackrel{\text{NLL}}{=} -K_H(\mu_H, \mu_{\mathcal{J}}) - K_J(\mu_J, \mu_{\mathcal{J}}) - K_S(\mu_S, \mu_{\mathcal{J}}),$$

$$\tilde{R}(e_\alpha, e_\beta) \stackrel{\text{NLL}}{=} \eta_J(\mu_J, \mu_{\mathcal{J}}) + \eta_S(\mu_S, \mu_{\mathcal{J}})$$

- Difference in the denominator:

(ignoring power-suppressed terms and terms beyond NLL)

<p style="color: orange;">Our result: <math>\Gamma(1 + \eta_J)\Gamma(1 + \eta_S)</math></p> <p>JHEP 1409 (2014) 046: <math>\Gamma(1 + \eta_J + \eta_S)</math></p>	}	<p>Difference at <math>\mathcal{O}(\alpha_s^2 \ln^2)</math></p> <p>in the bulk</p>
---	---	--

# Scale choices

- **Boundary conditions**

$$\Sigma(e_\alpha, e_\beta = e_\alpha^{\beta/\alpha}) = \Sigma(e_\alpha)$$

( $e_\beta$  has been integrated over its entire range)

$$\Sigma(e_\alpha = e_\beta, e_\beta) = \Sigma(e_\beta)$$

( $e_\alpha$  has been integrated over its entire range)


derivative:

$$\left. \frac{\partial}{\partial e_\alpha} \Sigma(e_\alpha, e_\beta) \right|_{e_\beta = e_\alpha^{\beta/\alpha}} = \frac{d\sigma}{de_\alpha}$$

$$\left. \frac{\partial}{\partial e_\alpha} \Sigma(e_\alpha, e_\beta) \right|_{e_\beta = e_\alpha} = 0$$

and similarly for  $\partial/\partial e_\beta$  with  $B1 \leftrightarrow B2$

- **Boundary conditions in JHEP 1409 (2014) 046 (Larkoski, Moult, Neill) fulfilled by adding power-suppressed**





# Scale choices

- **Boundary conditions**

$$\Sigma(e_\alpha, e_\beta = e_\alpha^{\beta/\alpha}) = \Sigma(e_\alpha)$$

( $e_\beta$  has been integrated over its entire range)

$$\Sigma(e_\alpha = e_\beta, e_\beta) = \Sigma(e_\beta)$$

( $e_\alpha$  has been integrated over its entire range)

derivative:

$$\left. \frac{\partial}{\partial e_\alpha} \Sigma(e_\alpha, e_\beta) \right|_{e_\beta = e_\alpha^{\beta/\alpha}} = \frac{d\Sigma}{de_\alpha}$$

$$\left. \frac{\partial}{\partial e_\alpha} \Sigma(e_\alpha, e_\beta) \right|_{e_\beta = e_\alpha} = 0$$

and similarly for  $\partial/\partial e_\beta$  with  $B1 \leftrightarrow B2$

- Boundary conditions in JHEP 1409 (2014) 046 (Larkoski, Moulton, Neill) fulfilled by adding power-suppressed

- **Profile scales**

Boundary conditions can be fulfilled by appropriate scale choice:

$$\mu_{\mathcal{S}}(e_\alpha, e_\beta) \Big|_{B1} = \mu_S(e_\alpha, e_\beta) \Big|_{B1}$$

$$\mu_{\mathcal{J}}(e_\alpha, e_\beta) \Big|_{B2} = \mu_J(e_\alpha, e_\beta) \Big|_{B2}$$

$$\left. \frac{\partial}{\partial e_\alpha} \mu_J(e_\alpha, e_\beta) \right|_{B2} = \frac{d}{de_\alpha} \mu_J(e_\alpha, e_\alpha^{\beta/\alpha})$$

$$\left. \frac{\partial}{\partial e_\alpha} \mu_{\mathcal{S}}(e_\alpha, e_\beta) \right|_{B2} = \frac{d}{de_\alpha} \mu_S(e_\alpha, e_\alpha^{\beta/\alpha})$$

$$\left. \frac{\partial}{\partial e_\alpha} \mu_S(e_\alpha, e_\beta) \right|_{B2} = \frac{d}{de_\alpha} \mu_S(e_\alpha, e_\alpha^{\beta/\alpha})$$

$$\left. \frac{\partial}{\partial e_\alpha} \mu_X(e_\alpha, e_\beta) \right|_{B1} = 0, \quad X = J, \mathcal{S}, S$$

and similarly for  $\partial/\partial e_\beta$

# Conclusions

- Resummation of double-differential measurements achieved via a new effective theory framework SCET+ containing collinear-soft modes
  - ↪ Factorization formula derived at the phase-space boundaries and in the intermediate regime. Continuous cross section description by matching factorization formula across different regions.
- Two applications we studied:
  - $pp \rightarrow Z + 0 \text{ jets}$ : jet veto is imposed through the beam thrust and transverse momentum of the  $Z$  measured
  - Measurement of two angularities on a single jet

**Thank you!**



**Back-up slides**

# Effective theory framework

- $n$ -collinear gauge transformation:

Groups together  $W_n^\dagger \xi_n$  ( $W_n^\dagger \rightarrow W_n^\dagger U_n^\dagger$ )

$$\xi_n \rightarrow U_n \xi_n, \quad W_n \rightarrow U_n W_n, \quad S_n \rightarrow S_n, \quad V_n \rightarrow V_n, \quad X_n \rightarrow X_n$$

Similarly  $\bar{\xi}_{\bar{n}} W_{\bar{n}}$  is grouped together by  $\bar{n}$ -collinear gauge transformation

- $n$ -collinear-soft gauge transformation:

Groups together  $V_n^\dagger X_n$

$$W_n^\dagger \xi_n \rightarrow W_n^\dagger \xi_n, \quad S_n \rightarrow S_n, \quad V_n \rightarrow U_{ncs} V_n, \quad X_n \rightarrow U_{ncs} X_n$$

Similarly  $X_{\bar{n}}^\dagger V_{\bar{n}}$  is grouped together by  $\bar{n}$ -collinear-soft gauge transformation

- soft gauge transformation:

$$\begin{aligned} W_n^\dagger \xi_n &\rightarrow W_n^\dagger \xi_n, & S_n &\rightarrow U_s S_n, & V_n &\rightarrow U_s V_n U_s^\dagger, & X_n &\rightarrow U_s X_n U_s^\dagger \\ \bar{\xi}_{\bar{n}} W_{\bar{n}} &\rightarrow \bar{\xi}_{\bar{n}} W_{\bar{n}}, & S_{\bar{n}} &\rightarrow U_s S_{\bar{n}}, & V_{\bar{n}} &\rightarrow U_s V_{\bar{n}} U_s^\dagger, & X_{\bar{n}} &\rightarrow U_s X_{\bar{n}} U_s^\dagger \end{aligned}$$

Fixes the remaining ordering

# Non-global logarithms

- To what extent can our framework be used to calculate non-global logarithms, arising when different restrictions are applied in different regions of phase space?
- Consider: Instead of measuring  $p_T$  of Z boson, measure  $p_T$  of ISR it recoils against (ISR in one hemisphere)

- Factorization theorem:

$$\begin{aligned} \frac{d^4\sigma}{dQ^2 dY dp_{T,\text{ISR}}^2 d\mathcal{T}} &= \sum_q \hat{\sigma}_q^0 H(Q^2, \mu) \int dt_2 \int d^2\vec{k}_{1\perp} d^2\vec{k}_{1\perp}^{\text{cs}} \int dk_1^+ dk^+ S(k^+, \mu) \\ &\times B_q(x_1, \vec{k}_{1\perp}, \mu, \nu) B_{\bar{q}}(t_2, x_2, \mu) \mathcal{S}(k_1^+, \vec{k}_{1\perp}^{\text{cs}}, \mu, \nu) \\ &\times \delta\left(\mathcal{T} - k_1^+ - \frac{e^Y t_2}{Q} - k^+\right) \delta\left(p_{T,\text{ISR}}^2 - |\vec{k}_{1\perp} + \vec{k}_{1\perp}^{\text{cs}}|^2\right) + (q \leftrightarrow \bar{q}) \end{aligned}$$

- This does not address the problem arising when the soft function contains multiple scales (e.g. when beam thus measurement would be restricted to one hemisphere)

# Matching of the effective theories

- At NNLL one can show:

$$\mathcal{I}_{qq}^{(1)}(t, x, \vec{k}_\perp) = \delta(t) \mathcal{I}_{qq}^{(1)}(x, \vec{k}_\perp) + \delta(1-x) \mathcal{S}^{(1)}(t/p^-, \vec{k}_\perp)$$

$$\mathcal{I}_{qg}^{(1)}(t, x, \vec{k}_\perp) = \delta(t) \mathcal{I}_{qg}^{(1)}(x, \vec{k}_\perp),$$

$$S^{(1)}(k^+, \vec{k}_\perp) = \frac{1}{\pi} \delta(\vec{k}_\perp^2) S^{(1)}(k^+) + 2\mathcal{S}^{(1)}(k^+, \vec{k}_\perp)$$

- Patch together the NNLL cross section

$$\begin{aligned} \frac{d^4\sigma}{dQ^2 dY dp_T^2 d\mathcal{T}} &= \sum_q \hat{\sigma}_q^0 H(Q^2) \int dt_1 dt_2 \int d^2\vec{k}_{1\perp} d^2\vec{k}_{2\perp} d^2\vec{k}_{1\perp}^{\text{cs}} d^2\vec{k}_{2\perp}^{\text{cs}} d^2\vec{k}_\perp \int dk_1^+ dk_2^+ dk^+ \\ &\times \left[ B_q(t_1, x_1, \vec{k}_{1\perp}) - \mathcal{S}^{(1)}(t_1 e^{-Y}/Q, \vec{k}_{1\perp}) \right] \mathcal{S}(k_1^+, \vec{k}_{1\perp}^{\text{cs}}) \\ &\times \left[ B_{\bar{q}}(t_2, x_2, \vec{k}_{2\perp}) - \mathcal{S}^{(1)}(t_2 e^Y/Q, \vec{k}_{2\perp}) \right] \mathcal{S}(k_2^+, \vec{k}_{2\perp}^{\text{cs}}) \\ &\times \left[ S(k^+, \vec{k}_\perp) - 2\mathcal{S}^{(1)}(k^+, \vec{k}_\perp) \right] \delta\left(\mathcal{T} - \frac{e^{-Y}t_1 + e^Y t_2}{Q} - k_1^+ - k_2^+ - k^+\right) \\ &\times \delta(p_T^2 - |\vec{k}_{1\perp} + \vec{k}_{2\perp} + \vec{k}_{1\perp}^{\text{cs}} + \vec{k}_{2\perp}^{\text{cs}} + \vec{k}_\perp|^2) + (q \leftrightarrow \bar{q}) \end{aligned}$$