Contribution ID: 112

Type: Working Group Sessions

Neutrino absorption in the Earth and measurement of the neutrino-nucleon cross-section at multi-TeV energies with IceCube

Tuesday, 20 June 2017 18:10 (20 minutes)

At the IceCube Neutrino Observatory, atmospheric and astrophysical neutrino fluxes become attenuated after passing through the Earth at energies above 40 TeV. This provides a unique method to measure the neutrino-nucleon cross section at energies significantly beyond previous accelerator-based approaches that reach 400 GeV. Here we report on the first measurement of neutrino absorption in the Earth, using a sample 10,784 energetic upward-going neutrino-induced muons observed in one year of IceCube data. From a fit to the two-dimensional distribution of muon energy and zenith angle, we measure the cross-section to be 1.30+0.21-0.19 (stat.) +0.39-0.43 (syst.) times the Standard Model expectation in an energy range from 6.3 TeV to 980 TeV.

Primary author: Dr MIARECKI, Sandra (United States Air Force Academy)

Co-authors: BINDER, Gary (Lawrence Berkeley National Laboratory/University of California Berkeley); Dr KLEIN, Spencer (LBNL & amp; UC Berkeley)

Presenter: BINDER, Gary (Lawrence Berkeley National Laboratory/University of California Berkeley)

Session Classification: Working Group: Astroparticle physics and cosmology

Track Classification: Astroparticle Physics and Cosmology Working Group