

The Short-Baseline Neutrino Program

Joseph Zennamo

On behalf of the MicroBooNE, ICARUS, and SBND Collaborations


48th Annual Fermilab Users Meeting
June 10th, 2015



THE UNIVERSITY OF
CHICAGO

Sterile Neutrinos

- Anomalies in a variety of experiments provide hints to the possible existence of sterile neutrinos
- The most significant of these come from LSND
- MiniBooNE followed up with additional anomalous results




These results can be interpreted as oscillations involving sterile neutrinos with $\Delta m^2 \sim 1\text{eV}^2$ and an $L/E \approx 1\text{ km/GeV}$

Sterile Neutrino Oscillations


- The large mass splitting means at short-baselines we are effectively decoupled from standard neutrino oscillations

$$P_{\nu_\alpha \rightarrow \nu_\beta}^{3+1} = \sin^2 2\theta \sin^2 \frac{\Delta m_{41}^2 L}{4E}$$


- We search for these oscillations by looking for the transformation of one active neutrino flavor into another
- Resolving these small effects requires control of systematic uncertainties




Booster Neutrino Beam




- Booster Neutrino Beam has been stably run for a decade and is well characterized
- 8 GeV Protons from the Booster collide with a beryllium target and the resulting mesons are focused by a magnetic horn
- This results is a beam of mostly muon neutrinos, where we can search for flavor disappearance and appearance




Short-Baseline Neutrino Program



Short-Baseline Neutrino Program



MicroBooNE
89t Active Mass




Short-Baseline Neutrino Program



SBND
112t Active Mass


- SBND will provide a detailed characterization of the beam before oscillations can occur
- This allows for the cancellation of many of the dominant systematics



Short-Baseline Neutrino Program




- The ICARUS T600 is the largest LAr TPC ever built, 7 times larger than MicroBooNE
- This mass provides us with high sensitivity to oscillated neutrinos allowing for a precision search



Short-Baseline Neutrino Program

- The goal of SBN is to perform the world's most sensitive sterile neutrino search
 - The three collaborations worked together to design such a program
- The resulting proposal contained a careful examination of all known backgrounds and systematic uncertainties

A Proposal for a Three Detector Short-Baseline Neutrino Oscillation Program in the Fermilab Booster Neutrino Beam




arXiv:1503.01520

“The Committee recommends Stage 1 approval for the SBN program...”

~Fermilab Physics Advisory Committee, January 2015



The Directorate has since granted Stage 1 approval!

Electron Neutrino Appearance





- The proposal leveraged the expertise gained by MiniBooNE both for the neutrino flux and the uncertainties on that flux
- By using a series of three LAr TPCs we are able to greatly reduce the photon-like backgrounds compared to MiniBooNE
- Given its large mass and far location ICARUS provides exquisite sensitivity to a potential oscillated signal




Sensitivity: Electron Neutrino Appearance



Disappearance Signature




Sensitivity: Muon Neutrino Disappearance




Ties to Long-Baseline Program

- This program is also a platform to develop the technical and analysis techniques necessary for DUNE
 - Bringing together the LAr TPC expertise from all over the world to collaborate on this program
 - Designing and validating event reconstruction on a large number of neutrino events and performing sensitive $\nu_\mu \rightarrow \nu_e$ oscillation analysis
 - Detailed studies of GeV-scale ν -Ar cross sections




Cross Sections at the SBN

- MicroBooNE will lay the ground work for an intense cross section program to characterize ν -Ar interactions
- Building off this the SBND will record **1.5 million** neutrino interactions **per year** allowing for detailed studies of rarer processes and expanded differential measurements
- The T600 sits in the NuMI beam far off-axis, allowing it to record a large statistics sample of higher energy neutrino interactions




Building the Program: SBND

- The near detector building is finalizing its design with construction scheduled to begin October 2015





- The designs for the cryostat and TPC are maturing quickly with close collaboration between Fermilab, CERN, and many institutions!



Building the Program: ICARUS T600

- The design of the far detector building has been finalized with construction starting this July!
- The refurbishment of the T600 has already begun at CERN
- The first of the two T300 modules will be finished by the end of 2015 and the second module by the end of 2016



Moving Forward

- The groundbreaking for the SBN detector buildings will take place this summer
- This starts an exciting period for all three collaborations, where activities range from design, to construction, to commissioning
- MicroBooNE will provide us the first look at what we might be seeing at short-baselines

**Finding Sterile Neutrinos Would be
Revolutionary!**

Thank You From

The ICARUS-WA104 Collaboration

M. Antonello¹⁶, B. Baibussinov³¹, V. Bellini⁵, P. Benetti³², S. Bertolucci⁶, H. Bilokon¹⁵, F. Boffelli³², M. Bonesini¹⁷, J. Bremer⁶, E. Calligarich³², S. Centro³¹, A.G. Cocco¹⁹, A. Dermenev²⁰, A. Falcone³², C. Farnese³¹, A. Fava³¹, A. Ferrari⁶, D. Gibin³¹, S. Gninenko²⁰, N. Golubev²⁰, A. Guglielmi³¹, A. Ivashkin²⁰, M. Kirsanov²⁰, J. Kisiel³⁸, U. Kose⁶, F. Mammoliti⁵, G. Mannocchi¹⁵, A. Menegolli³², G. Meng³¹, D. Mladenov⁶, C. Montanari³², M. Nessi⁶, M. Nicoletto³¹, F. Noto⁶, P. Picchi¹⁵, F. Pietropaolo³¹, P. Płoński⁴², R. Potenza⁵, A. Rappoldi³², G. L. Raselli³², M. Rossella³², C. Rubbia^{*6,11,16}, P. Sala¹⁸, A. Scaramelli¹⁸, J. Sobczyk⁴⁴, M. Spanu³², D. Stefan¹⁸, R. Sulej⁴³, C.M. Sutera⁵, M. Torti³², F. Tortorici⁵, F. Varanini³¹, S. Ventura³¹, C. Vignoli¹⁶, T. Wachala¹², and A. Zani³²

The LAr1-ND Collaboration

C. Adams⁴⁵, C. Andreopoulos²³, A. Ankowski⁴¹, J. Asaadi⁴⁰, L. Bagby¹⁰, B. Baller¹⁰, N. Barros³³, M. Bass³⁰, S. Bertolucci⁶, M. Bishai³, A. Bitadze²⁵, J. Bremer⁶, L. Bugel²⁶, L. Camilleri⁹, F. Cavanna^{a,10}, H. Chen³, C. Chi⁹, E. Church¹⁰, D. Cianci⁷, G. Collin²⁶, J.M. Conrad²⁶, G. De Geronimo³, R. Dharmapalan¹, Z. Djurcic¹, A. Ereditato², J. Esquivel⁴⁰, J. Evans²⁵, B.T. Fleming⁴⁵, W.M. Foreman⁷, J. Freestone²⁵, T. Gamble³⁷, G. Garvey²⁴, V. Genty⁹, D. Göldi², H. Greenlee¹⁰, R. Guenette³⁰, A. Hackenburg⁴⁵, R. Hänni², J. Ho⁷, J. Howell¹⁰, C. James¹⁰, C.M. Jen⁴¹, B.J.P. Jones²⁶, L.M. Kalousis⁴¹, G. Karagiorgi²⁵, W. Ketchum²⁴, J. Klein³³, J. Klinger³⁷, U. Kose⁶, I. Kreslo², V.A. Kudryavtsev³⁷, D. Lissauer³, P. Livesly²², W.C. Louis²⁴, M. Lu□thi², C. Mariani⁴¹, K. Mavrokoridis²³, N. McCauley²³, N. McConkey³⁷, I. Mercer²², T. Miao¹⁰, G.B. Mills²⁴, D. Mladenov⁶, D. Montanari¹⁰, J. Moon²⁶, Z. Moss²⁶, S. Mufson¹⁴, M. Nessi⁶, B. Norris¹⁰, F. Noto⁶, J. Nowak²², S. Pal³⁷, O. Palamara^{*b,10}, J. Pater²⁵, Z. Pavlovic¹⁰, J. Perkin³⁷, G. Pulliam⁴⁰, X. Qian³, L. Qiuguang²⁴, V. Radeka³, R. Rameika¹⁰, P.N. Ratoff²², M. Richardson³⁷, C. Rudolf von Rohr², D.W. Schmitz^{*7}, M.H. Shaevitz⁹, B. Sippach⁹, M. Soderberg⁴⁰, S. Söldner-Rembold²⁵, J. Spitz²⁶, N. Spooner³⁷, T. Strauss², A.M. Szczec^{25,45}, C.E. Taylor²⁴, K. Terao⁹, M. Thiesse³⁷, L. Thompson³⁷, M. Thomson⁴, C. Thorn³, M. Toups²⁶, C. Touramanis²³, R.G. Van De Water²⁴, M. Weber², D. Whittington¹⁴, T. Wongjirad²⁶, B. Yu³, G.P. Zeller¹⁰, and J. Zennamo⁷

The MicroBooNE Collaboration



R. Acciarri¹⁰, C. Adams⁴⁵, R. An¹³, A. Ankowski⁴¹, J. Asaadi⁴⁰, L. Bagby¹⁰, B. Baller¹⁰, G. Barr³⁰, M. Bass³⁰, M. Bishai³, A. Blake⁴, T. Bolton²¹, C. Bromberg²⁷, L. Bugel²⁶, L. Camilleri⁹, D. Caratelli⁹, B. Carls¹⁰, F. Cavanna^{a,10}, H. Chen³, E. Church¹⁰, G.H. Collin²⁶, J.M. Conrad²⁶, M. Convery³⁹, S. Dytmar³⁴, B. Eberly³⁹, A. Ereditato², J. Esquivel⁴⁰, B.T. Fleming^{*45}, W.M. Foreman⁷, V. Genty⁹, D. Göldi², S. Gollapinni²¹, M. Graham³⁹, E. Gramellini⁴⁵, H. Greenlee¹⁰, R. Grossi⁸, R. Guenette³⁰, A. Hackenburg⁴⁵, O. Hen²⁶, J. Hewes²⁵, J. Ho⁷, G. Horton-Smith²¹, C. James¹⁰, C.M. Jen⁴¹, R.A. Johnson⁸, B.J.P. Jones²⁶, J. Joshi³, H. Jostlein¹⁰, D. Kaleko⁹, L. Kalousis⁴¹, G. Karagiorgi²⁵, W. Ketchum²⁴, B. Kirby³, M. Kirby¹⁰, T. Kobilarcik¹⁰, I. Kreslo², Y. Li³, B. Littlejohn¹³, D. Lissauer³, S. Lockwitz¹⁰, W.C. Louis²⁴, M. Lu□thi², B. Lundberg¹⁰, A. Marchionni¹⁰, C. Mariani⁴¹, J. Marshall⁴, K. McDonald³⁵, V. Meddage²¹, T. Miceli²⁸, G.B. Mills²⁴, J. Moon²⁶, M. Mooney³, M.H. Moulaï²⁶, R. Murrells²⁵, D. Naples³⁴, P. Nienaber³⁶, O. Palamara^{b,10}, V. Paolone³⁴, V. Papavassiliou²⁸, S. Pate²⁸, Z. Pavlovic¹⁰, S. Pordes¹⁰, G. Pulliam⁴⁰, X. Qian³, J.L. Raaf¹⁰, V. Radeka³, R. Rameika¹⁰, B. Rebel¹⁰, L. Rochester³⁹, C. Rudolf von Rohr², B. Russell⁴⁵, D.W. Schmitz⁷, A. Schukraft¹⁰, W. Seligman⁹, M. Shaevitz⁹, M. Soderberg⁴⁰, J. Spitz²⁶, J. St. John⁸, T. Strauss², A.M. Szczec^{25,45}, N. Tagg²⁹, K. Terao⁹, M. Thomson⁴, C. Thorn³, M. Toups²⁶, Y. Tsai³⁹, T. Usher³⁹, R. Van de Water²⁴, M. Weber², S. Wolbers¹⁰, T. Wongjirad²⁶, K. Woodruff²⁸, M. Xu¹³, T. Yang¹⁰, B. Yu³, G.P. Zeller^{*10}, J. Zennamo⁷, and C. Zhang³

Additional Fermilab Contributors


W. Badgett¹⁰, K. Biery¹⁰, S. Brice¹⁰, S. Dixon¹⁰, M. Geynisman¹⁰, C. Moore¹⁰, E. Snider¹⁰, and P. Wilson¹⁰

Backups

Effect of More Statistics





SBN Detectors



Systematics



Systematic Uncertainties: Neutrino Flux

- Hadron Production
 - Charged Pions
 - Kaons
- Beam Focusing
 - Current
 - Skin Effects
- Secondary Interactions
 - Within the target and horn







Systematics

Systematic Uncertainties: Neutrino Flux




Systematics



Systematics



Systematic Uncertainties: ν -Argon Cross Sections

The three detectors use the same target nucleus but differences in the flux, detector acceptances, and other effects can lead to a different composition of event types which can lead to systematic differences between the detectors







Systematics

Systematic Uncertainties: ν -Argon Cross Sections




Systematics



Systematics

Systematic Uncertainties: “Dirt” Background




Systematics

Systematic Uncertainties: Cosmic Ray Background

- One positive aspect of cosmic induced background is that we can measure it directly with arbitrarily high precision
- Using off beam random triggers we will be able to build a large sample of cosmic events
 - The systematic uncertainty on this background becomes the statistical uncertainty of those measurements

ArgonTube Cosmic Ray Events



Muon Neutrino Flux at NDOS

