

Success of the International Collaboration

James Stone

Boston University

Some thoughts for LBNE.

Slides 10th Anniversary of Super-Kamiokande

Oct. 3, 2012

The 1980's ...

- Both IMB and Kamiokande Collaborations published many papers on the same topics.
- A friendly scientific competition spanned a decade.
- Independent scientific results made a strong impact on the physics community.

Search for Proton Decay into $e^+ \pi^0$

R. M. Bionta, G. Blewitt, C. B. Bratton, B. G. Cortez,^(a) S. Errede, G. W. Forster,^(a) W. Gajewski, M. Goldhaber, J. Greenberg, T. J. Haines, T. W. Jones, D. Kielczewska,^(b) W. R. Kropp, J. G. Learned, E. Lehmann, J. M. LoSecco, P. V. Ramana Murthy,^(c) H. S. Park, F. Reines, J. Schultz, E. Shumard, D. Sinclair, D. W. Smith,^(d) H. W. Sobel, J. L. Stone, L. R. Sulak, R. Svoboda, J. C. van der Velde, and C. Wuest

The University of California at Irvine, Irvine, California 92717, and The University of Michigan, Ann Arbor, Michigan 48109, and Brookhaven National Laboratory, Upton, New York 11973, and California Institute of Technology, Pasadena, California 91125, and Cleveland State University, Cleveland, Ohio 44115, and The University of Hawaii, Honolulu, Hawaii 96822, and University College, London WC1E 6BT, United Kingdom

(Received 13 April 1983)

Observations were made 1570 meters of water equivalent underground with an 8000-metric-ton water Cherenkov detector. During a live time of 80 d no events consistent with the decay $p \rightarrow e^+ \pi^0$ were found in that the limit on the lifetime for bou ratio is $\tau/B > 6.5 \times 10^{31}$ yr; for free dence). Observed cosmic-ray muon

PACS numbers: 13.30.Eg, 11.30.Ly

Journal of the Physical Society of Japan
Vol. 54, No. 9, September, 1985, pp. 3213-3216

LETTERS

Search for Nucleon Decay into Charged Lepton + Mesons

Katsushi ARISAKA, Takaaki KAJITA, Masatoshi KOSHIBA,
Masayuki NAKAHATA, Yuichi OYAMA, Atsuto SUZUKI,
Masato TAKITA, Yoji TOTSUKA, Tadashi KIFUNE,[†]
Teruhiro SUDA,[†] Kasuke TAKAHASHI^{††} and Kazumasa MIYANO^{†††}

Department of Physics and ICEPP, University of Tokyo, Tokyo 113

[†]*Institute for Cosmic Ray Research, University of Tokyo, Tokyo 188*

^{††}*KEK, National Laboratory for High Energy Physics, Ibaraki 305*

^{†††}*Department of Physics, University of Niigata, Niigata 950-21*

(Received July 19, 1985)

With a 3000 ton water Cerenkov detector operated 2700 m.w.e. underground, 103 fully contained events were observed during a live time of 343 days. Most of the events are well interpreted as due to ν interactions. Four multi-ring events survive after applying criteria for nucleon decay. The lower limits on τ/B obtained from these data exceed 10^{31} yr (90% C.L.) for most of the possible decay modes.

IMB

Proton Decay

Kamiokande

SU(5)

Search for High-Energy Muons from Cygnus X-3

Y. Oyama, K. Arisaka,^(a) T. Kajita, M. Koshiba, M. Nakahata, A. Suzuki, M. Takita, and Y. Totsuka

*Department of Physics, and International Center for Elementary Particle Physics,
University of Tokyo, Tokyo 113, Japan*

T. Kifune and T. Suda

Institute for Cosmic Ray Research, University of Tokyo, Tokyo 188, Japan

N. Sato

Department of Astronomy, University of Tokyo, Tokyo 113, Japan

K. Takahashi

National Laboratory for High Energy Physics (KEK), Ibaraki 305, Japan

and

K. Miyano

Department of Physics, University of Niigata, Niigata 950-21, Japan

(Received 15 October 1985)

A total of 1.9×10^6 good-quality high-energy muon events were analyzed from the data of the underground Kamioka nucleon-decay experiment. No definite sign of excess flux was observed in the direction of Cygnus X-3 and in its phase interval of 0.7 to 0.8. The 90%-C.L. (confidence level) upper limit for the excess flux is $2.2 \times 10^{-12} \text{ cm}^{-2} \text{ sec}^{-1}$ for rock overburden greater than 2400 m of water equivalent.

Kamiokande

Cygnus X-3

IMB

Underground search for muons correlated with Cygnus X-3

R. M. Bionta,¹ G. Blewitt,^d C. B. Bratton,^e D. Casper,^{b,n} A. Ciocio,ⁿ R. Claus,ⁿ
M. Crouch,¹ S. T. Dye,^f S. Errede,¹ G. W. Foster,^o W. Gajewski,^a K. S. Ganezer,^a
M. Goldhaber,^c T. J. Haines,^a T. W. Jones,^b D. Kielczewska,^b W. R. Kropp,^a
J. G. Learned,^f J. M. LoSecco,^m J. Matthews,^b H. S. Park,^k L. R. Price,^a F. Reines,^a
J. Schulz,^a S. Seidel,^{b,n} E. Shumard,^p D. Sinclair,^b H. W. Sobel,^a J. L. Stone,ⁿ L. Sulak,ⁿ
R. Svoboda,^a G. Thornton,^b J. C. van der Velde,^b and C. Wuest¹

^aThe University of California, Irvine, California 92717

^bThe University of Michigan, Ann Arbor, Michigan 48109

^cBrookhaven National Laboratory, Upton, New York 11973

^dCalifornia Institute of Technology, Pasadena, California 91125

^eCleveland State University, Cleveland, Ohio 44115

^fThe University of Hawaii, Honolulu, Hawaii 96822

^gUniversity College, London WC1E 8BT, United Kingdom

^hWarsaw University, Warsaw PL-00-681, Poland

ⁱCase Western Reserve, Cleveland Ohio 44106

^jThe University of Illinois, Urbana, Illinois 61801

^kUniversity of California, Berkeley, California 94720

^lLawrence Livermore National Laboratory, Livermore, California 94550

^mNotre Dame University, Notre Dame, Indiana 46556

ⁿBoston University, Boston, Massachusetts 02215

Observation of a Neutrino Burst from the Supernova SN1987A

K. Hirata,^(a) T. Kajita,^(a) M. Koshiba,^(a,b) M. Nakahata,^(b) Y. Oyama,^(b)
 N. Sato,^(c) A. Suzuki,^(b) M. Takita,^(b) and Y. Totsuka^(a,c)
University of Tokyo, Tokyo 113, Japan

T. Kifune and T. Suda

Institute for Cosmic Ray Research, University of Tokyo, Tokyo 118, Japan

K. Takahashi and T. Tanimori

National Laboratory for High Energy Physics (KEK), Ibaraki 305, Japan

K. Miyano and M. Yamada

Department of Physics, University of Niigata, Niigata 950-21, Japan

E. W. Beier, L. R. Feldscher, S. B. Kim, A. K. Mann, F. M. Newcomer, R. Van Berg, and W. Zhang
Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania 19104

and

B. G. Cortez^(d)

California Institute of Technology, Pasadena, California 91125

(Received 10 March 1987)

A neutrino burst was observed in the Kamiokande II detector on 23 February 1987, 7:35:35 UT (± 1 min) during a time interval of 13 sec. The signal consisted of eleven electron events of energy 7.5 to 36 MeV, of which the first two point back to the Large Magellanic Cloud with angles $18^\circ \pm 18^\circ$ and $15^\circ \pm 27^\circ$.

Kamiokande

Supernova 1987A

IMB

Observation of a Neutrino Burst in Coincidence with Supernova 1987A in the Large Magellanic Cloud

R. M. Bionta,⁽¹²⁾ G. Blewitt,⁽⁴⁾ C. B. Bratton,⁽⁵⁾ D. Casper,^(2,14) A. Ciocio,⁽¹⁴⁾ R. Claus,⁽¹⁴⁾ B. Cortez,⁽¹⁶⁾
 M. Crouch,⁽⁹⁾ S. T. Dye,⁽⁶⁾ S. Errede,⁽¹⁰⁾ G. W. Foster,⁽¹⁵⁾ W. Gajewski,⁽¹⁾ K. S. Ganezer,⁽¹⁾
 M. Goldhaber,⁽³⁾ T. J. Haines,⁽¹⁾ T. W. Jones,⁽⁷⁾ D. Kielczewska,^(1,8) W. R. Kropp,⁽¹⁾ J. G. Learned,⁽⁶⁾
 J. M. LoSecco,⁽¹³⁾ J. Matthews,⁽²⁾ R. Miller,⁽¹⁾ M. S. Mudan,⁽⁷⁾ H. S. Park,⁽¹¹⁾ L. R. Price,⁽¹⁾
 F. Reines,⁽¹⁾ J. Schultz,⁽¹⁾ S. Seidel,^(2,14) E. Shumard,⁽¹⁶⁾ D. Sinclair,⁽²⁾ H. W. Sobel,⁽¹⁾ J. L. Stone,⁽¹⁴⁾
 L. R. Sulak,⁽¹⁴⁾ R. Svoboda,⁽¹⁾ G. Thornton,⁽²⁾ J. C. van der Velde,⁽²⁾ and C. Wuest⁽¹²⁾

⁽¹⁾The University of California, Irvine, Irvine, California 92717

⁽²⁾The University of Michigan, Ann Arbor, Michigan 48109

⁽³⁾Brookhaven National Laboratory, Upton, New York 11973

⁽⁴⁾California Institute of Technology, Jet Propulsion Laboratory, Pasadena, California 91109

⁽⁵⁾Cleveland State University, Cleveland, Ohio 44115

⁽⁶⁾The University of Hawaii, Honolulu, Hawaii 96822

⁽⁷⁾University College, London WC1E 6BT, United Kingdom

⁽⁸⁾Warsaw University, Warsaw, Poland

⁽⁹⁾Case Western Reserve University, Cleveland, Ohio 44106

⁽¹⁰⁾The University of Illinois, Urbana, Illinois 61801

⁽¹¹⁾The University of California, Berkeley, California 94720

⁽¹²⁾Lawrence Livermore National Laboratory, Livermore, California 94550

⁽¹³⁾The University of Notre Dame, Notre Dame, Indiana 46556

⁽¹⁴⁾Boston University, Boston, Massachusetts 02215

⁽¹⁵⁾Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁽¹⁶⁾AT&T Bell Laboratories, Summit, New Jersey 07910

(Received 13 March 1987)

A burst of eight neutrino events preceding the optical detection of the supernova in the Large Magellanic Cloud has been observed in a large underground water Cherenkov detector. The events span an interval of 6 s and have visible energies in the range 20–40 MeV.

The Early 1990's ...

- Low energy threshold becomes important for supernovae neutrinos and solar neutrinos.
- IMB's small PMTs not competitive for this physics.
- Easter weekend, 1991, IMB tank developed a leak that caused much damage to the detector. IMB Finished.

INSTITUTE FOR COSMIC RAY RESEARCH

ICRR

UNIVERSITY OF TOKYO

Now funded!

ICRR-Report-227-90-20

SUPER-KAMIOKANDE

Y. Totsuka

(Dec. 1990)

Invited talk presented at the International Symposium
on Underground Physics Experiments, at Science
Council of Japan, April, 1990

3-2-1 Midori-cho Tanashi, Tokyo 188 Japan

Telephone (0424)-61-4131, Telefax (0424)-68-1438

First Contact

Email sent to Professor Totsuka on January 23, 1992 follows a conference phone call the previous night.

IMB proposed:

- Merge forces on Super-Kamiokande
- Use IMB PMTs to build anti-detector (outer)
- Independent simulations
- Work on KEK beam test
- Visit Japan
- Totsuka to visit Boston

```
#1 23-JAN-1992 15:55:26.85 NEWMAIL
\address
Professor Y. Totsuka
Institute for Cosmic Ray Research
University of Tokyo
5-1-1 Midori-Cho Tanashi
Tokyo 188, Japan
\body
Dear Professor Totsuka:
\baselineskip=15pt
We were very pleased to talk with you today about the future plans of
Super-Kamiokande and the future possibilities of IMB.
Let me take this opportunity to reiterate to you, on behalf of the IMB
collaboration, the possible collaboration that has been discussed at our
recent IMB group meeting. We are truly impressed with the promise
presented by your proposed Super-Kamiokande detector. This detector
offers a natural extension of the beautiful results from Kamiokande,
particularly the proton decay limits, the solar neutrino observations
Press RETURN for more...
MAIL>
#1 23-JAN-1992 15:55:26.85 NEWMAIL
and the discovery of supernova neutrinos. Although the IMB
collaboration has also had the pleasure of setting significant limits on
proton decay, astrophysical point sources, monopoles, wimps, etc. as
well as observing the neutrinos from the supernova, we recognize the
superior capabilities of Super-Kamiokande. Seven times the fiducial
volume of IMB would extend our reach in the search for proton decay, etc
way beyond our current hopes. What you have done so well already in
Kamiokande clearly will be much better done at Super-Kamiokande and we
wonder if merging forces to develop together the world's largest
detector might even hold more promise.
At this point in the evolution of IMB, considerable resources could be
brought to a new collaborative effort. We have in hand 2000
characterized, well-working phototubes housed for high pressure water
operation. High voltage, electronics and readout are all completely
operational. These, for example, could be put to service either at a
test beam or as the active veto for your large detector.
Super-Kamiokande will not be duplicated. As you know, we have developed
Press RETURN for more...
MAIL>
#1 23-JAN-1992 15:55:26.85 NEWMAIL
a detailed knowledge of the very same technology that you use. Two
independent simulations and double checks of detector performance could
be extremely valuable. In IMB, we currently lack a crucial Monte Carlo
ingredient, a detailed neutrino calibration at an accelerator.
Therefore, we look eagerly toward your proposed study of  $e\bar{\nu}_e$  and  $\tau$  neutrino interactions at KEK. Since we are limited by our
knowledge of the neutrino interactions, your topology and cross section
measurements, particularly for multi-ring events, are a project that we
would dearly like to participate in.
MAIL>
#1 23-JAN-1992 15:55:26.85 NEWMAIL
April 29. We would like to extend an invitation to you to come to Boston
at that time and see our technical and analysis facilities. We would be
pleased to arrange for your travel from Philadelphia or whatever may be
appropriate. For the IMB Collaboration, we look forward to discussing this
with you. Thank you very much for taking the time to discuss these
prospects with us today.
\closing
Sincerely,
\vskip 1cm
S.T. Dye
Associate Professor
\vskip 1cm
J.L. Stone
Professor
```

Response from Professor Totsuka

Excerpted from letter dated:

2/17/92

The summary of our discussions are;

(1) We are in general quite positive to organize an international collaboration for the Super-Kamiokande experiment, especially to collaborate with the IMB group, which we greatly respect for their excellent achievements on proton decay and neutrino physics.

(2) Obviously the collaborating people must agree on the importance of the physics goals, and the design of the Super-Kamiokande detector.

(3) It is indeed a very good idea to have the anti-counters from the beginning by help of the foreign participation. In this respect we welcome your proposal of bringing the IMB inventories to Kamioka.

(4) Of course there are many technical details that have to be worked out. Therefore, we welcome some of you from Boston and Irvine to come to the ICRR, Tokyo and to see the Kamioka facility. We like to discuss the matters that were already addressed in your conversation with Yoichiro Suzuki.


(5) Also we would like to discuss with you on the initial beam-test experiment (with e and μ beams) scheduled in 1993 (construction in 1992).

(6) Sometime later when we are agreeable on the technical points, we would like to discuss the issues related to

(i) a possible contract between your institute(s) and ours.

(ii) a structure of the collaboration, such as the role of the collaboration meeting, etc.

Sincerely yours



Yoji Totsuka

The First Celebration ...

At ICRR Tanashi Campus in spring of 1992, our agreement to work together on Super-Kamiokande and to prepare a formal collaboration document is celebrated.



Collaboration Agreement Signed October 18, 1992 in Takayama

SuperKamiokande Collaboration Agreement Between the Collaborating Groups

Purpose

The purpose of this document is to define the terms and conditions under which the collaborating groups (at present Japanese and American) agree to work together in building and operating a 50,000 ton water Cherenkov detector at the Kamioka mine. The experiment shall be known as SuperKamiokande. The goals of the experiment include a search for nucleon decay, atmospheric neutrino studies, solar neutrinos, and studies of/searches for other astrophysical and particle physics phenomena. It is agreed that all collaborating groups are free to participate in all aspects of the experiment.

History

The SuperKamiokande (SK) project was initially considered and later materialized in a detailed form by M. Koshiba and his collaborators in 1984. After many years of intense effort, the full project was funded by the Japanese Ministry of Education, Science, and Culture in 1991. The final design and production of various detector parts started the same year.

In February 1992, some members of the IMB Collaboration called the SK Spokesman (Y. Totsuka) and expressed an interest in joining the SK Collaboration.

Several meetings and discussions have followed the initial February 1992 contact by the IMB group. This agreement is the result of our efforts to merge the two largest water Cherenkov groups in the world for the purpose of building and operating SuperKamiokande.

Organization of the Collaboration

The SK Collaboration has 4 levels of organization:

- 1) Spokesman (Yoji Totsuka)
- 2) Executive Committee (Initially, Y. Totsuka, K. Nakamura, A. Suzuki, H. Sobel, and J. Stone)
- 3) Collaboration Council (Defined below)
- 4) Collaboration Meeting (All Collaborators)

Signatures

Yoji Totsuka 10/18/92
Date

Yoji Totsuka
Institute for Cosmic Ray Research - University of Tokyo
Spokesman for the SuperKamiokande Collaboration

Atsuto Suzuki 10/18/92
Date

Atsuto Suzuki
KEK

Kenzo Nakamura 10/18/92
Date

Kenzo Nakamura
Institute of Cosmic Ray Research
University of Tokyo

Members of the Executive Committee

Henry W. Sobel 10/18/92
Date

Henry Sobel
University of California - Irvine

James Stone 18 Oct 92
Date

James Stone
Boston University

Members of the Executive Committee

M. Koshiba Oct. 18, 92
Date

Masatoshi Koshiba
Tokai University

John Learned 18
Date

John Learned
University of Hawaii

Teruhiro Suda Oct 20, 92
Date

Teruhiro Suda
Kobe University

Todd Haines 20 Oct 92
Date

Todd Haines
University of Maryland

Kazumasa Miyano 10/18/92
Date

Kazumasa Miyano
Niigata University

Robert Svoboda 19 Nov. 92
Date

Robert Svoboda
Louisiana State University

Yorikazu Nagashima 10/18/92
Date

Yorikazu Nagashima
Osaka University

Clyde B. Bratton 20 Nov
Date

Clyde B. Bratton
Cleveland State University

The first project performed together as a Collaboration was the KEK Beam Test

ca. 1993 - 95

This test served as a detailed calibration of the 50 cm diameter PMTs used in Kamiokande and 8 inch PMTs used in the IMB Experiments.

