

Heavy Flavour Physics

Lecture 2 of 2

Tim Gershon
University of Warwick

HCPSS 2010

Fifth CERN-Fermilab Hadron Collider Physics Summer School


19th August 2010



Contents

- Tuesday
 - What is “heavy flavour physics”?
 - Why is it interesting?
 - What do we know about it as of today?
- Today
 - What do we hope to learn from current and future heavy flavour experiments?


Summary from Tuesday



Adding a few other constraints we find

$$\bar{\rho} = 0.132 \pm 0.020$$

$$\bar{\eta} = 0.358 \pm 0.012$$



Consistent with Standard Model fit

- some “tensions”


Still plenty of room for new physics

Topics to cover today

- Flavour physics at hadron colliders (mainly LHCb)
- More on CP violation
 - The third Unitarity Triangle angle: γ
 - Tree-dominated decays vs. loop-dominated decays
 - CP violating phase in B_s^0 oscillations
 - CP violating phase in D^0 oscillations
- Rare decays
 - $B_s^0 \rightarrow \mu\mu$, $B \rightarrow K^*\mu\mu$, $B_s^0 \rightarrow \phi\gamma$
- Future experiments


Flavour physics at hadron colliders

	$e^+e^- \rightarrow \Upsilon(4s) \rightarrow B\bar{B}$ PEP-II, KEK-B	$p\bar{p} \rightarrow b\bar{b}X$ ($\sqrt{s} = 2$ TeV) TeVatron	$pp \rightarrow b\bar{b}X$ ($\sqrt{s} = 14$ TeV) LHC
prod	1 nb	$\sim 100 \mu\text{b}$	$\sim 500 \mu\text{b}$
typ. $b\bar{b}$ rate	10 Hz	~ 100 kHz	~ 500 kHz
purity	$\sim 1/4$	$\sigma_{b\bar{b}}/\sigma_{inel} \approx 0.2\%$	$\sigma_{b\bar{b}}/\sigma_{inel} \approx 0.6\%$
pile-up	0	1.7	0.5-20
B content	B^+B^- (50%), $B^0\bar{B}^0$ (50%)	B^+ (40%), B^0 (40%), B_s (10%), B_c ($< 1\%$), b-baryons (10%)	
B boost	small, $\beta\gamma \sim 0.56$	large, decay vertices are displaced	
event structure	BB pair alone	many particles non-associated to $b\bar{b}$	
prod. vertex	Not reconstructed	reconstructed with many tracks	
$B^0\bar{B}^0$ mixing	coherent	incoherent \rightarrow flavour tagging dilution	



Geometry

- In high energy collisions, $b\bar{b}$ pairs produced predominantly in forward or backward directions
- LHCb is a forward spectrometer



The LHCb Detector
JINST 3 (2008) S08005






LHCb detector features

- Tracking and calorimetry
 - basic essentials of any collider experiment!
 - muon chambers
- VELO
 - reconstruct displaced vertices
- RICH
 - particle identification (K/ π separation)
- Trigger
 - fast and efficient

More details in dedicated
detector lectures


Status of LHCb at end 2009

~ 1/nb




Status of LHCb early 2010


~ 10/nb




π^0




η



ϕ





D



J/psi


B



VELO: best resolution of any LHC vertex detector

Status of LHCb at ICHEP2010

~ 100/nb



LHCb yield


60 events / 180/nb
 (consistent with MC based expectation)

Compare CDF (CDF note 10071)
 45000 events / 4.3/fb

Compare D0 (PRL 100 (2008) 211802)
 40000 events / 2.8/fb

Compare Belle (arXiv:1008.2567)
 41000 events / 711/fb



Still a long way to go but catching up fast!



Already > 1/pb on tape

Measurement of the $b\bar{b}$ cross-section at LHCb

- Use $D^0 \rightarrow K^- \pi^+$ decays associated with a μ^-
- Separate “prompt” from “D from B” events with a fit to the impact parameter distribution
- Perform analysis in bins of η
 - LHCb acceptance: $2 < \eta < 6$



$$\sigma(pp \rightarrow b\bar{b}X; 2 < \eta < 6) = (74.9 \pm 5.3 \pm 12.8) \mu\text{b}$$

Use PYTHIA to extrapolate to full range
 $\sigma(pp \rightarrow b\bar{b}X) = (282 \pm 20 \pm 48) \mu\text{b}$

largest systematics from luminosity, tracking efficiency

S.Stone and A.Golutvin at ICHEP 2010

It's all about the trigger

Challenge is


- to efficiently select most interesting B decays
- while maintaining manageable data rates

Main backgrounds

- “minimum bias” inelastic pp scattering
- other charm and beauty decays

Handles

- high p_T signals (muons)
- displaced vertices



Spectroscopy

- I've talked about the headline items of flavour physics
 - CP violation, searches for new physics
 - what we tell the funding agencies, and the press
- But, much of the physics performed by flavour experiments is the study of properties of hadronic states
 - lifetimes, masses, decay channels, quantum numbers
 - and the discoveries of new ones

1) Observation of a narrow meson decaying to $D^+(s)\pi^0$ at a mass of $2.32\text{-GeV}/c^2$.
By BABAR Collaboration (Bernard Aubert *et al.*). SLAC-PUB-9711, BABAR-PUB-03-011, Apr 2003. 7pp.
[Press Release from SLAC.](#)
Published in *Phys.Rev.Lett.***90:242001,2003.**
e-Print: [hep-ex/0304021](#)

TOPCITE = 500+

[References](#) | [LaTeX\(US\)](#) | [LaTeX\(EU\)](#) | [Harvmac](#) | [BibTeX](#) | [Keywords](#) | Cited [521 times](#)
[Abstract](#) and [Postscript](#) and [PDF](#) from arXiv.org (mirrors: [au](#) [br](#) [cn](#) [de](#) [es](#) [fr](#) [il](#) [in](#) [it](#) [jp](#) [kr](#) [ru](#) [tw](#) [uk](#) [za](#) [aps](#) [lanl](#))
Journal Server [doi:[10.1103/PhysRevLett.90.242001](#)]
[BaBar Publications Database](#)
[BaBar Password Protected Publications Database](#)
[CERN Library Record](#)
[pdgLive \(measurements quoted by PDG\)](#)
[Press Release about this paper](#)
[SLAC Document Server](#)
[EXP SLAC-PEP2-BABAR](#)
[Bookmarkable link to this information](#)

Observation of a narrow charmonium - like state in exclusive $B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$ decays.
By Belle Collaboration (S.K. Choi *et al.*). Sep 2003. 10pp.
[Press release.](#)
Published in *Phys.Rev.Lett.***91:262001,2003.**
e-Print: [hep-ex/0309032](#)

TOPCITE = 500+

[References](#) | [LaTeX\(US\)](#) | [LaTeX\(EU\)](#) | [Harvmac](#) | [BibTeX](#) | [Keywords](#) | Cited [514 times](#)
[Abstract](#) and [Postscript](#) and [PDF](#) from arXiv.org (mirrors: [au](#) [br](#) [cn](#) [de](#) [es](#) [fr](#) [il](#) [in](#) [it](#) [jp](#) [kr](#) [ru](#) [tw](#) [uk](#) [za](#) [aps](#) [lanl](#))
Journal Server [doi:[10.1103/PhysRevLett.91.262001](#)]
[pdgLive \(measurements quoted by PDG\)](#)
[Press Release about this paper](#)
[EXP KEK-BF-BELLE](#)
[Bookmarkable link to this information](#)

Most highly cited papers from BaBar and Belle

Discovery of the lightest $b\bar{b}$ state – 2008

Observation of the Bottomonium Ground State in the Decay $Y(3S) \rightarrow \gamma \eta_b$



B. Aubert,¹ M. Bona,¹ Y. I. Karasik,¹ J. R. Loy,¹ V. Pojma,¹ E. Preprint,¹ X. Rodas,¹ V. Tisserand,¹ J. Garra Tico,² F. Graess,² L. Lopez,³ A. P. ...
The BaBar Collaboration Abrams,⁵ M. Rattalora,⁵

- Only recoil γ is reconstructed


$$m(\eta_b(1S)) = (9388.9^{+3.1}_{-2.3} \pm 2.7) \text{ MeV}/c^2$$

$$m(Y(1S)) - m(\eta_b(1S)) = (71.4^{+2.3}_{-3.1} \pm 2.7) \text{ MeV}/c^2$$

$$B(Y(3S) \rightarrow \gamma \eta_b(1S)) = (4.8 \pm 0.5 \pm 1.2) \times 10^{-4}$$



→ subtract smoothly varying background



Why wasn't the η_b discovered at a hadronic experiment?

- Remember: $Y(1S)$ discovered at FNAL in 1977
 - fixed target experiment: p on Be
- η_b is lighter
- Hadron collisions produce all types of b hadrons
- So why couldn't the η_b be discovered, e.g., at the Tevatron?

PRL 39 (1977) 252

Why wasn't the η_b discovered at a hadronic experiment?

- Remember: $Y(1S)$ discovered at FNAL in 1977
 - fixed target experiment: p on Be
- η_b is lighter
- Hadron collisions produce all types of b hadrons
- So why couldn't the η_b be discovered, e.g., at the Tevatron?
- **It's all about the trigger!**
 - need clean signature for trigger and reconstruction
 - CDF search used $\eta_b \rightarrow J/\psi J/\psi$ decay, with predicted BF $\sim 0!$

PRL 39 (1977) 252

Digression on a digression: The “Oops Leon”

Observation of High-Mass Dilepton Pairs in Hadron Collisions at 400 GeV

D. C. Hom, L. M. Lederman, H. P. Paar, H. D. Snyder, J. M. Weiss, and J. K. Yoh
Columbia University, New York, New York 10027*

and

J. A. Appel, B. C. Brown, C. N. Brown, W. R. Innes, and T. Yamanouchi
Fermi National Accelerator Laboratory, Batavia, Illinois 60510†

and

D. M. Kaplan
State University of New York at Stony Brook, Stony Brook, New York 11794*
(Received 28 January 1976)

We report preliminary results on the production of electron-positron pairs in the mass range 2.5 to 20 GeV in 400-GeV p -Be interactions. 27 high-mass events are observed in the mass range 5.5–10.0 GeV corresponding to $\sigma = (1.2 \pm 0.5) \times 10^{-35}$ cm² per nucleon. Clustering of 12 of these events between 5.8 and 6.2 GeV suggests that the data contain a new resonance at 6 GeV.

PRL 36 (1976) 1236

Homework exercise:

1. Read this paper
2. Do you find the “discovery” convincing?
3. Explain what's wrong





FIG. 2. Electron-positron mass spectrum: $d\sigma/dm$ per nucleon versus the effective mass. A linear $A\sigma$ -dependence is assumed. Note bin-width changes.

b hadron spectroscopy – Observation of the Ω_b^-

CDF PRD 80 (2009) 72003





$$m(\Omega_b^-) =$$

$$6054.4 \pm 6.8 \text{ (stat.)} \pm 0.9 \text{ (syst.) MeV}$$

$$6165 \pm 10 \text{ (stat)} \pm 13 \text{ (syst.) MeV}$$


significant discrepancy
to be understood

D0 PRL 101 (2008) 232002




b hadron spectroscopy – Observation of the Σ_b

CDF PRL 99 (2007) 202001




Fully hadronic decay chain:



Impressive demonstration of
B physics potential with
hadronic triggers


More b hadron spectroscopy

Study of the quantum numbers of X(3872)



PRL 98 (2007) 132002

Discovery of the Y(4140) in $B \rightarrow J/\psi\phi K$




CDF Note 10244 & PRL 102 (2009) 242002

The smoking gun exotic hadron: A charged charmonium-like state

$$B^0 \rightarrow Z(4430)^- K^+, Z(4430)^- \rightarrow \psi' \pi^-$$

Belle PRL 100 (2008) 142001




Clear peak

Still there in more detailed analysis

PRD 80 (2009) 031104

BABAR PRD 79 (2009) 112001



Data consistent with $K\pi$ reflections

Slight peak but no evidence for new state

But also consistent with Belle

Need more experimental input (CDF, D0 or LHCb)

OK, back to weak physics

Direct CP violation

- Condition for DCPV: $|\bar{A}/A| \neq 1$
- Need \bar{A} and A to consist of (at least) two parts
 - with different weak (φ) and strong (δ) phases
- Often realised by “tree” and “penguin” diagrams


$$A = |T|e^{i(\delta_T - \phi_T)} + |P|e^{i(\delta_P - \phi_P)} \quad \bar{A} = |T|e^{i(\delta_T + \phi_T)} + |P|e^{i(\delta_P + \phi_P)}$$

$$A_{CP} = \frac{|\bar{A}|^2 - |A|^2}{|\bar{A}|^2 + |A|^2} = \frac{2|T||P|\sin(\delta_T - \delta_P)\sin(\phi_T - \phi_P)}{|T|^2 + |P|^2 + 2|T||P|\cos(\delta_T - \delta_P)\cos(\phi_T - \phi_P)}$$

Homework:
prove it

Example: $B \rightarrow K\pi$

(weak phase difference is γ)



Feynman tree (a) and penguin (b) diagrams for the $B_d^0 \rightarrow K^+ \pi^-$ decay

The famous penguin story

Penguin diagram

From Wikipedia, the free encyclopedia

In quantum field theory, **penguin diagrams** are a class of Feynman diagrams which are important for understanding CP violating processes in the standard model.

They were first isolated and studied by Mikhail Shifman, Arkady Vainshtein, and Valentin Zakharov.^[1] The processes which they describe were first directly observed in 1991 and 1994 by the CLEO collaboration.

Origin of the name


[edit]

John Ellis was the first to refer to a certain class of Feynman diagrams as **penguin diagrams**, due in part to their shape, and in part to a legendary bar-room bet with Melissa Franklin. According to John Ellis:^[2]

“ Mary K. [Gaillard], Dimitri [Nanopoulos] and I first got interested in what are now called penguin diagrams while we were studying CP violation in the Standard Model in 1976... The penguin name came in 1977, as follows.

In the spring of 1977, Mike Chanowitz, Mary K and I wrote a paper on GUTs predicting the b quark mass before it was found. When it was found a few weeks later, Mary K, Dimitri, Serge Rudaz and I immediately started working on its phenomenology. That summer, there was a student at CERN, Melissa Franklin who is now an experimentalist at Harvard. One evening, she, I, and Serge went to a pub, and she and I started a game of darts. We made a bet that if I lost I had to put the word penguin into my next paper. She actually left the darts game before the end, and was replaced by Serge, who beat me. Nevertheless, I felt obligated to carry out the conditions of the bet.

For some time, it was not clear to me how to get the word into this b quark paper that we were writing at the time. Then, one evening, after working at CERN, I stopped on my way back to my apartment to visit some friends living in Meyrin where I smoked some illegal substance. Later, when I got back to my apartment and continued working on our paper, I had a sudden flash that the famous diagrams look like penguins. So we put the name into our paper, and the rest, as they say, is history.



The famous penguin story

Penguin diagram

From Wikipedia, the free encyclopedia

In quantum field theory, **penguin diagrams** are a class of Feynman diagrams which are important for understanding CP violating processes in the standard model.

They were first isolated and studied by ... describe were first directly observed in 1991 and 1994 by the CLEO collaboration



Origin of the name

John Ellis was the first to refer to a ... shape, and in part to a legendary bar-ro

“ Mary K. [Gaillard], Dimitri [... penguin diagrams while we ... penguin name came in 197


In the spring of 1977, Mik ... quark mass before it was fo ... Rudaz and I immediately st ... student at CERN, Melissa ... she, I, and Serge went to a ... lost I had to put the word p ... the end, and was replaced b ... conditions of the bet.

For some time, it was not ... writing at the time. Then, one evening, after working at CERN, I stopped on my way back to my apartment to visit some friends living in Meyrin where I smoked some illegal substance. Later, when I got back to my apartment and continued working on our paper, I had a sudden flash that the famous diagrams look like penguins. So we put the name into our paper, and the rest, as they say, is history.



Example of a penguin diagram

Direct CP asymmetries in charmless hadronic B decays



Direct CP violation in $B \rightarrow K\pi$

- Direct CP violation in $B \rightarrow K\pi$ sensitive to γ
 too many hadronic parameters \Rightarrow need theory input
 NB. interesting deviation from naïve expectation

“K π puzzle”

$$A_{CP}(K^- \pi^+) = (-9.8_{-1.1}^{+1.2})\% \quad A_{CP}(K^- \pi^0) = (5.0 \pm 2.5)\%$$


$$\Delta(A_{CP}) = (-14.8 \pm 2.8)\%$$

HFAG averages

BABAR PRD 76 (2007) 091102 & arXiv:0807.4226; also CDF

Could be a sign of new physics ...
 ... first need to rule out possibility of larger
 than expected QCD corrections

Belle Nature 452 (2008) 332



Clean observables in $B \rightarrow K\pi$ (etc.)

- Measure more $B_{u,d} \rightarrow K\pi$ decays & relate by isospin
- Perform similar analysis on $B \rightarrow K^*\pi$ &/or $B \rightarrow K\rho$
 - Dalitz plot analyses of $K\pi\pi$ final states extract both amplitudes and relative phases \rightarrow more observables
- Measure $B_s \rightarrow KK$ decays & relate by U-spin
 - e.g. relation between time-dependent CP violation observables in $B_s \rightarrow K^+K^-$ and $B^0 \rightarrow \pi^+\pi^-$
- Dalitz plot analyses of $B_s \rightarrow KK\pi$

Note: flavour symmetries very useful

But, still get theory error from symmetry breaking (difficult to evaluate)


... data driven methods will win in the end (unless miracle breakthrough)

$B \rightarrow h^+ h'^-$ at hadron colliders


- Excellent channel if you can trigger on it (displaced vertex)
- Particle ID extremely important

- Key channel for LHCb

first time-dependent measurements ASAP



A. Golutvin at ICHEP 2010



CDF PRL 103 (2009) 031801. See also D.Tonelli at Beauty 2009

Importance of γ from $B \rightarrow DK$



- γ plays a unique role in flavour physics

the only CP violating parameter that can be measured through tree decays (*)

(*) more-or-less

- A benchmark Standard Model reference point
 - doubly important after New Physics is observed

$$\propto V_{cb} V_{us}^*$$




$$\propto V_{ub} V_{cs}^*$$

Variants use different B or D decays
require a final state common to both D^0 and \bar{D}^0

Why is $B \rightarrow DK$ so nice?

- For theorists:
 - theoretically clean: no penguins; factorisation works
 - all parameters can be determined from data
- For experimentalists:
 - many different observables (different final states)
 - all parameters can be determined from data
 - γ & δ_B (weak & strong phase differences), r_B (ratio of amplitudes)







B → DK methods

- Different D decay final states
 - CP eigenstates, e.g. K^+K^- (GLW)
 - doubly-Cabibbo-suppressed decays, e.g. $K^+\pi^-$ (ADS)
 - singly-Cabibbo-suppressed decays, e.g., $K^{*+}K^-$ (GLS)
 - self-conjugate multibody decays, e.g., $K_S \pi^+\pi^-$ (GGSZ)
- Different B decays
 - $B^- \rightarrow DK^-, D^*K^-, DK^{*-}$ never studied before (or not much)
 - $B^0 \rightarrow DK^{*0}$ (or $B \rightarrow DK\pi$ Dalitz plot analysis)
 - $B^0 \rightarrow DK_S, B_S^0 \rightarrow D\phi$ (with or without time-dependence)
 - $B_S^0 \rightarrow D_S K, B^0 \rightarrow D^{(*)}\pi$ (time-dependent)

Latest results on $B \rightarrow DK$: GLW


Evidence for direct CP violation ($\gamma \neq 0$)



BABAR arXiv:1007.0504


A_{CP} Averages

HFAG
ICHEP 2010
PRELIMINARY





Latest results on $B \rightarrow DK$: GGSZ

BABAR arXiv:1005.1096



BELLE PRD 81 (2010) 112002



$$\gamma = (68^{+15}_{-14} \pm 4 \pm 3)^\circ$$

Uncertainty due to assumed $D \rightarrow K_S \pi^+ \pi^-$ decay model

$$\Phi_3 = (78^{+11}_{-12} \pm 4 \pm 9)^\circ$$

Evidence for direct CP violation ($\gamma \neq 0$)

Model independent $B \rightarrow DK$ Dalitz measurements

- Use CP-tagged CLEOc data to measure average $D^0-\bar{D}^0$ phase difference

CLEO-c Results: c_i & s_i

NEW

A.Powell at Beauty 2009




[Phys Rev. D 80, 032002 (2009)]

- Result \pm stat \pm sys \pm ($K_L\pi\pi$ $K_S\pi\pi$ syst)

i	c_i	s_i
1	$0.743 \pm 0.037 \pm 0.022 \pm 0.013$	$0.014 \pm 0.160 \pm 0.077 \pm 0.045$
2	$0.611 \pm 0.071 \pm 0.037 \pm 0.009$	$0.014 \pm 0.215 \pm 0.055 \pm 0.017$
3	$0.059 \pm 0.063 \pm 0.031 \pm 0.057$	$0.609 \pm 0.190 \pm 0.076 \pm 0.037$
4	$-0.495 \pm 0.101 \pm 0.052 \pm 0.045$	$0.151 \pm 0.217 \pm 0.069 \pm 0.048$
5	$-0.911 \pm 0.049 \pm 0.032 \pm 0.021$	$-0.050 \pm 0.183 \pm 0.045 \pm 0.036$
6	$-0.736 \pm 0.066 \pm 0.030 \pm 0.018$	$-0.340 \pm 0.187 \pm 0.052 \pm 0.047$
7	$0.157 \pm 0.074 \pm 0.042 \pm 0.051$	$-0.827 \pm 0.185 \pm 0.060 \pm 0.036$
8	$0.403 \pm 0.046 \pm 0.021 \pm 0.002$	$-0.409 \pm 0.158 \pm 0.050 \pm 0.002$

- Statistical uncertainties dominant
- c_i better determined than s_i
- Results also available for c_i' & s_i'
- Broad agreement with model predictions

- γ Uncertainty: $\sigma_{\text{CLEO-input}}(\gamma) = 1.7^\circ$
(recall model error = 7°)



The other Unitarity Triangles

- High statistics available at LHCb will allow sensitivity to smaller CP violating effects
 - CP violating phase in B_s oscillations ($O(\lambda^4)$)
 - B_s oscillations (Δm_s) measured 2006 (CDF)
 - CP violating phase in D^0 oscillations ($O(\lambda^5)$)
 - D^0 oscillations ($x_D = \Delta m_D / \Gamma_D$ & $y_D = 2\Delta\Gamma_D / \Gamma_D$) measured 2007 (Babar, Belle, later CDF)
- Observations of CP violation in both K^0 and B^0 systems won Nobel prizes!

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates

$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} + \mathcal{A}_{\text{CP}}^{\text{dir}} \cos(\Delta m t) + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} + \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 + a) e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} - \mathcal{A}_{\text{CP}}^{\text{dir}} \cos(\Delta m t) + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} - \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right].$$

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates

$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} + \mathcal{A}_{CP}^{dir} \cos(\Delta m t) + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} + \mathcal{A}_{CP}^{mix} \sin(\Delta m t) \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 - a) e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} - \mathcal{A}_{CP}^{dir} \cos(\Delta m t) + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} - \mathcal{A}_{CP}^{mix} \sin(\Delta m t) \right].$$

CP violating asymmetries

CP conserving parameter

$$A_{CP}^{dir} = C_{CP} = \frac{1 - |\lambda_{CP}|^2}{1 + |\lambda_{CP}|^2} \quad A_{\Delta\Gamma} = \frac{2 \Re(\lambda_{CP})}{1 + |\lambda_{CP}|^2} \quad A_{CP}^{mix} = S_{CP} = \frac{2 \Im(\lambda_{CP})}{1 + |\lambda_{CP}|^2}$$

$$(A_{CP}^{dir})^2 + (A_{\Delta\Gamma})^2 + (A_{CP}^{mix})^2 = 1$$

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates

$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} \text{ () } + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} \text{ ()} \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 + a) e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} \text{ () } + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} \text{ ()} \right].$$

- Untagged analyses still sensitive to some interesting physics

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates

$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} + 0 + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} + \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 + 0) e^{-\Gamma t} \times \left[\cosh \frac{\Delta\Gamma t}{2} - 0 + \mathcal{A}_{\Delta\Gamma} \sinh \frac{\Delta\Gamma t}{2} - \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right].$$

- In some channels, expect no direct CP violation
- and/or no CP violation in mixing

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates

$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\mathbf{1} + \mathcal{A}_{\text{CP}}^{\text{dir}} \cos(\Delta m t) + \mathbf{0} + \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 + a) e^{-\Gamma t} \times \left[\mathbf{1} - \mathcal{A}_{\text{CP}}^{\text{dir}} \cos(\Delta m t) + \mathbf{0} - \mathcal{A}_{\text{CP}}^{\text{mix}} \sin(\Delta m t) \right].$$

- In some channels, expect no direct CP violation
- B_d case: $\Delta\Gamma$ negligible

Time-dependent CP Violation Formalism

- Generic (but shown for B_s) decays to CP eigenstates


$$\Gamma(B_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} e^{-\Gamma t} \times \left[\mathbf{1} + \mathcal{A}_{\text{CP}}^{\text{dir}} \mathbf{1} + \mathcal{A}_{\Delta\Gamma} y\Gamma t + \mathcal{A}_{\text{CP}}^{\text{mix}} x\Gamma t \right]$$

$$\Gamma(\bar{B}_s(t) \rightarrow f) = \mathcal{N}_f |A_f|^2 \frac{1 + |\lambda_f|^2}{2} (1 + a) e^{-\Gamma t} \times \left[\mathbf{1} - \mathcal{A}_{\text{CP}}^{\text{dir}} \mathbf{1} + \mathcal{A}_{\Delta\Gamma} y\Gamma t - \mathcal{A}_{\text{CP}}^{\text{mix}} x\Gamma t \right].$$


- In some channels, expect no direct CP violation
- B_d case: $\Delta\Gamma$ negligible
- D^0 case: both $x = \Delta m/\Gamma$ and $y = \Delta\Gamma/2\Gamma$ small

Charm mixing and CP violation

HFAG world average Including results from BABAR, Belle, CDF, CLEO(c), FOCUS



Inconsistent with no mixing point (0,0)




Consistent with no CP violation point (1,0)

At LHCb can use $D \rightarrow K^+K^-$ to measure

- $A_{\Delta\Gamma} y_D$ (untagged or tagged); $A_{CP}^{\text{mix}} x_D$ (tagged)

Many other possible channels

$\beta_s (B_s \rightarrow J/\psi\phi)$



• VV final state

three helicity amplitudes

→ mixture of CP-even and CP-odd

disentangled using angular & time-dependent distributions

→ additional sensitivity

many correlated variables

→ complicated analysis

- 2004 K.Gibson at Beauty 2009
 - CDF: measurement of $\Delta\Gamma/\Gamma$
 - D0 measurement in 2005
- 2006
 - D0: first untagged analysis for Φ_s
 - CDF analysis in 2007
 - CDF: first measurement of Δm_s
- 2007
 - CDF: first flavour tagged analysis
 - D0 measurement in 2008
- 2008/9
 - First attempts at averages → → →
official CDF/D0 combination
 - Updated results ... both now 2.8/fb
- 2010
 - CDF & D0 results updated (5–6/fb)

$B_s \rightarrow J/\psi\phi$ formalism

Differential decay rate:

$$\frac{d^4\Gamma(B_s^0 \rightarrow J/\psi\phi)}{dt d\cos\theta d\varphi d\cos\psi} \equiv \frac{d^4\Gamma}{dt d\Omega} \propto \sum_{k=1}^6 h_k(t) f_k(\Omega)$$

B_s

\bar{B}_s

k	$h_k(t)$	$h_k(t)$	$f_k(\theta, \psi, \varphi)$
1	$ A_0(t) ^2$	$ \bar{A}_0(t) ^2$	$2\cos^2\psi(1 - \sin^2\theta\cos^2\varphi)$
2	$ A_{\parallel}(t) ^2$	$ \bar{A}_{\parallel}(t) ^2$	$\sin^2\psi(1 - \sin^2\theta\sin^2\varphi)$
3	$ A_{\perp}(t) ^2$	$ \bar{A}_{\perp}(t) ^2$	$\sin^2\psi\sin^2\theta$
4	$\Im\{A_{\parallel}^*(t)A_{\perp}(t)\}$	$\Im\{\bar{A}_{\parallel}^*(t)\bar{A}_{\perp}(t)\}$	$-\sin^2\psi\sin 2\theta\sin\varphi$
5	$\Re\{A_0^*(t)A_{\parallel}(t)\}$	$\Re\{\bar{A}_0^*(t)\bar{A}_{\parallel}(t)\}$	$\frac{1}{\sqrt{2}}\sin 2\psi\sin^2\theta\sin 2\varphi$
6	$\Im\{A_0^*(t)A_{\perp}(t)\}$	$\Im\{\bar{A}_0^*(t)\bar{A}_{\perp}(t)\}$	$\frac{1}{\sqrt{2}}\sin 2\psi\sin 2\theta\cos\varphi$

$A_0(0) \rightarrow$ CP even
 $A_{\parallel}(0) \rightarrow$ CP even
 $A_{\perp}(0) \rightarrow$ CP odd

Derivations left as an exercise for the student (joke)

\pm signs differ for B_s and \bar{B}_s

$$|\bar{A}_0(t)|^2 = |\bar{A}_0(0)|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin\Phi \sin(\Delta m_s t) \right],$$

$$|\bar{A}_{\parallel}(t)|^2 = |\bar{A}_{\parallel}(0)|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin\Phi \sin(\Delta m_s t) \right],$$

$$|\bar{A}_{\perp}(t)|^2 = |\bar{A}_{\perp}(0)|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) + \sin\Phi \sin(\Delta m_s t) \right],$$

$$\Im\{\bar{A}_{\parallel}^*(t)\bar{A}_{\perp}(t)\} = |\bar{A}_{\parallel}(0)||\bar{A}_{\perp}(0)| e^{-\Gamma_s t} \left[-\cos(\delta_{\perp} - \delta_{\parallel}) \sin\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin(\delta_{\perp} - \delta_{\parallel}) \cos(\Delta m_s t) + \cos(\delta_{\perp} - \delta_{\parallel}) \cos\Phi \sin(\Delta m_s t) \right],$$

$$\Re\{\bar{A}_0^*(t)\bar{A}_{\parallel}(t)\} = |\bar{A}_0(0)||\bar{A}_{\parallel}(0)| e^{-\Gamma_s t} \cos\delta_{\parallel} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin\Phi \sin(\Delta m_s t) \right] \text{ and}$$


$$\Im\{\bar{A}_0^*(t)\bar{A}_{\perp}(t)\} = |\bar{A}_0(0)||\bar{A}_{\perp}(0)| e^{-\Gamma_s t} \left[-\cos\delta_{\perp} \sin\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin\delta_{\perp} \cos(\Delta m_s t) + \cos\delta_{\perp} \cos\Phi \sin(\Delta m_s t) \right].$$

$\beta_s (B_s \rightarrow J/\psi\phi)$

G.Punzi at EPS 2009


- Tevatron measurements using tagged $B_s \rightarrow J/\psi\phi$
- Angular analyses of vector-vector final state
- Results depend on $\Delta\Gamma$

CDF note 9787



$3166 \pm 56 B_s \rightarrow J/\psi\phi$ events $\beta_s^{J/\psi\phi}$ [rad]

D0 5928-CONF




$1967 \pm 65 B_s \rightarrow J/\psi\phi$ events $\beta_s^{J/\psi\phi}$ [rad]

$\beta_s (B_s \rightarrow J/\psi\phi)$

- Teva
- Angu
- Resu

CD




68% CL interval: Φ_s in $[0.27, 0.59] \cup [0.97, 1.30]$ rad

95% CL interval: Φ_s in $[0.10, 1.42]$ rad

$\beta_s (B_s \rightarrow J/\psi\phi)$ – LHCb prospects

MC performance:

- 50k events / fb^{-1} consistent with number of $B_s \rightarrow J/\psi\phi$ candidates seen in data
- $\langle\sigma_t\rangle = 0.038$ ps. Present resolution in data is ~ 1.6 worse but sufficient for $\Delta m_s \sim 17.7/\text{ps}$ (adds 30% dilution to the sensitivity)
- Tagging performance $\varepsilon D^2 = 6.2\%$ will be tested with more data




A.Golutvin at ICHEP2010

Rare Decays

$b \rightarrow s\gamma$ rate and photon energy spectrum

Archetypal FCNC probe for new physics




Belle PRL 103 (2009) 241801

$$B(B \rightarrow X_s \gamma)_{E_\gamma > 1.7 \text{ GeV}} = (3.45 \pm 0.15 \pm 0.40) \times 10^{-4}$$

consistent with the SM prediction


$b \rightarrow s\gamma$ photon polarisation measurement

- Search for time-dependent asymmetry
- Observable effect requires NP: left-handed current & new CP phase



Excellent prospects for LHCb with $B_s \rightarrow \phi\gamma$


Belle PRL 100 (2008) 121801



Can also use, eg., $B \rightarrow K^*e^+e^-$ (low q^2)

$$B \rightarrow K^* \mu^+ \mu^-$$

- $b \rightarrow s l^+ l^-$ processes also governed by FCNCs
 - rates and asymmetries of many exclusive processes sensitive to NP
- Queen among them is $B_d \rightarrow K^{*0} \mu^+ \mu^-$
 - superb laboratory for NP tests
 - **experimentally clean signature**
 - many kinematic variables ...
 - ... with clean theoretical predictions (at least at low q^2)



Operator Product Expansion

Build an effective theory for b physics

- take the weak part of the SM
- integrate out the heavy fields (W,Z,t)
- (like a modern version of Fermi theory for weak interactions)

$$\mathcal{L}_{(\text{full EW}\times\text{QCD})} \longrightarrow \mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{QED}\times\text{QCD}} \left(\begin{array}{l} \text{quarks } \neq t \\ \& \text{ leptons} \end{array} \right) + \sum_n C_n(\mu) Q_n$$

Q_n – local interaction terms (operators), C_n – coupling constants (Wilson coefficients)

Wilson coefficients

- encode information on the weak scale
- are calculable and known in the SM (at least to leading order)
- are affected by new physics

For $K^*\mu\mu$ we care about C_7 (also affects $b \rightarrow s\gamma$), C_9 and C_{10}

Effective operators

$$\mathcal{H}_W^{\Delta B=1, \Delta C=0, \Delta S=-1} = 4 \frac{G_F}{\sqrt{2}} \left(\lambda_c^s (C_1(\mu) Q_1^c(\mu) + C_2(\mu) Q_2^c(\mu)) \right. \\ \left. + \lambda_u^s (C_1(\mu) Q_1^u(\mu) + C_2(\mu) Q_2^u(\mu)) - \lambda_t^s \sum_{i=3}^{10} C_i(\mu) Q_i(\mu) \right)$$

where the $\lambda_q^s = V_{qb}^* V_{qs}$ and the operator basis is given by

$$\begin{aligned} Q_1^q &= \bar{b}_L^\alpha \gamma^\mu q_L^\alpha \bar{q}_L^\beta \gamma_\mu s_L^\beta & Q_2^q &= \bar{b}_L^\alpha \gamma^\mu q_L^\beta \bar{q}_L^\beta \gamma_\mu s_L^\alpha \\ Q_3 &= \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q \bar{q}_L^\beta \gamma_\mu q_L^\beta & Q_4 &= \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q \bar{q}_L^\beta \gamma_\mu q_L^\alpha \\ Q_5 &= \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q \bar{q}_R^\beta \gamma_\mu q_R^\beta & Q_6 &= \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q \bar{q}_R^\beta \gamma_\mu q_R^\alpha \\ Q_7 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q e_q \bar{q}_R^\beta \gamma_\mu q_R^\beta & Q_8 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q e_q \bar{q}_R^\beta \gamma_\mu q_R^\alpha \\ Q_9 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q e_q \bar{q}_L^\beta \gamma_\mu q_L^\beta & Q_{10} &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q e_q \bar{q}_L^\beta \gamma_\mu q_L^\alpha \end{aligned}$$

Four-fermion operators (except $Q_{7\gamma}$ & Q_{8g}) – dimension 6

$$Q_{7\gamma} = \frac{e}{16\pi^2} m_b \bar{b}_L^\alpha \sigma^{\mu\nu} F_{\mu\nu} s_L^\alpha$$

$$Q_{8g} = \frac{g_s}{16\pi^2} m_b \bar{b}_L^\alpha \sigma^{\mu\nu} G_{\mu\nu}^A T^A s_L^\alpha$$


$$Q_{9V} = \frac{1}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \bar{l} \gamma_\mu l$$

$$Q_{10A} = \frac{1}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \bar{l} \gamma_\mu \gamma_5 l$$

Theory of $B \rightarrow K^* \mu^+ \mu^-$

- Given for inclusive $b \rightarrow s \mu^+ \mu^-$ for simplicity
 - physics of exclusive modes \approx same but equations are more complicated (involving form factors, etc.)
- Differential decay distribution

$$\frac{d^2\Gamma}{dq^2 d \cos \theta_l} = \frac{3}{8} \left[(1 + \cos^2 \theta_l) H_T(q^2) + 2 \cos \theta_l H_A(q^2) + 2 (1 - \cos^2 \theta_l) H_L(q^2) \right]$$



$$H_T(q^2) \propto 2q^2 \left[\left(C_9 + 2C_7 \frac{m_b^2}{q^2} \right)^2 + C_{10}^2 \right],$$


$$H_A(q^2) \propto -4q^2 C_{10} \left(C_9 + 2C_7 \frac{m_b^2}{q^2} \right),$$

$$H_L(q^2) \propto \left[(C_9 + 2C_7)^2 + C_{10}^2 \right].$$

This term gives a forward-backward asymmetry

Results on $B \rightarrow K^* \mu^+ \mu^-$ kinematic distributions

- Differential decay rate
- Longitudinal polarisation
- Forward-backward asymmetry




CDF note 10047
See also Babar PRD 79 (2009) 031102(R)
and Belle PRL 103 (2009) 171801

SM prediction of the
zero crossing point is
particularly clean

$$B_s \rightarrow \mu^+ \mu^-$$

Killer app. for new physics discovery

- Very small in the SM
- Huge NP enhancement
($\tan \beta = \text{ratio of Higgs vevs}$)
- Clean experimental signature



$$BR(B_s \rightarrow \mu^+ \mu^-)^{SM} = (3.3 \pm 0.3) \times 10^{-8} \quad BR(B_s \rightarrow \mu^+ \mu^-)^{MSSM} \propto \tan^6 \beta / M_{A^0}^4$$

$B_s \rightarrow \mu\mu$ – Latest Results






$$B(B_s \rightarrow \mu\mu) < 5.1 \times 10^{-8} \text{ @95\% CL}$$

$$B(B_s \rightarrow \mu\mu) < 4.3 \times 10^{-8} \text{ @95\% CL}$$

$$B(B_d \rightarrow \mu\mu) < 7.6 \times 10^{-9} \text{ @95\% CL}$$

D0 arXiv:1006.3469

CDF Public Note 9892




$B_s \rightarrow \mu\mu$ – LHC Prospects

- Not only LHCb!
- GPD discovery potential
- LHCb should have advantage
 - but let's wait and see ...

Mass resolutions (MeV/c²)

	ATLAS	CMS	LHCb
$B_s \rightarrow \mu\mu$	90	36	18
$B_s \rightarrow D_s\pi$	53		14
$B_s \rightarrow J/\psi\phi$	61	14	16

V. Gibson HCPSS 2008



Expect ~ 10 SM events in 1/fb

LHCb: A. Golutvin at ICHEP 2010

LHCb upgrade

- To fully exploit LHC potential for heavy flavour physics will require an upgrade to LHCb
 - full readout & trigger at 40 MHz to enable high L running
 - “high L” = $10^{34}/\text{cm}^2/\text{s}$ (so independent of machine upgrade)
 - planned for 2016 shutdown

LHC machine schedule



Other future flavour experiments


- SuperKEKB & Belle2 (upgraded KEKB & Belle)
- SuperB (new e^+e^- facility proposed in Italy)
 - $B \rightarrow \tau \nu$, inclusive measurements, τ physics, ...
- Rare kaon decays
 - $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ (NA62, CERN); $K^0 \rightarrow \pi^0 \nu \bar{\nu}$ (KOTO, J-PARC)
- Muon to electron conversion (charged lepton flavour violation)
 - COMET/PRIME (J-PARC); mu2e (FNAL)

B → τν and charged Higgs limits


- Pure leptonic decays of charged B mesons very clean
 - clean SM prediction
 - clean effect of charged Higgs (2HDM or SUSY)

$$BR(B^+ \rightarrow l^+ \nu)^{SM} = \frac{G_F m_B}{8\pi} m_l^2 \left(1 - \frac{m_l^2}{m_B^2}\right)^2 f_B^2 |V_{ub}|^2 \tau_B \quad BR(B^+ \rightarrow l^+ \nu)^{NP} = BR(B^+ \rightarrow l^+ \nu)^{SM} \left(1 - \frac{m_B^2}{m_H^2} \tan^2 \beta\right)^2$$

Derivations left as exercise for the student (partial joke)



Existing (2009) flavour constraints rule out ATLAS discovery with 30/fb! (but there's always wiggle room)



Summary

- We still don't know:
 - why there are so many fermions in the SM
 - what causes the baryon asymmetry of the Universe
 - where exactly the new physics is ...
 - ... and what its flavour structure is
- Prospects are good for progress in the next few years
- We need a continuing programme of flavour physics into the 2020s
 - complementary to the high- p_T programme of the LHC

References and background reading

- Reviews by the Particle Data Group
 - <http://pdg.lbl.gov/>
- Heavy Flavour Averaging Group (HFAG)
 - <http://www.slac.stanford.edu/xorg/hfag/>
- CKMfitter & UTfit
 - <http://ckmfitter.in2p3.fr/> & <http://www.utfit.org/>
- Review journals (e.g. Ann. Rev. Nucl. Part. Phys.)
 - <http://nucl.annualreviews.org>
 - arXiv:0907.5386 (CKM 2008 write-up, to appear Phys. Repts.)
- Books
 - CP violation, I.I.Bigi and A.I.Sanda (CUP)
 - CP violation, G.C.Branco, L.Lavoura & J.P.Silva (OUP)

CKM

2010

<http://ckm2010.warwick.ac.uk>