

Search for Proton Medium Modifications in the $^4\text{He}(\text{e},\text{e}'\text{p})$ Reaction

Steffen Strauch
University of South Carolina

NuInt07: Fifth International Workshop on Neutrino-Nucleus Interactions in the Few-GeV Region
May 30, 2007 - June 3, 2007, Fermilab, Batavia, Illinois USA

Motivation

- Accurate description of $A(e,e')$ data sets strong restrictions on nuclear models of neutrino reactions, $A(\nu_l, l^-)$
- Nuclear reaction-model ingredients:
 - ▶ Relativity
 - ▶ Final-state interactions
 - ▶ Medium modifications
 - ▶ ...
- Exclusive $A(e,e'p)$ data provide for additional, sensitive tests
 - ▶ Cross sections
 - ▶ Polarization observables

e.g., O. Benhar *et al.*, Phys. Rev. D **72**, 053005 (2005); J.E. Amaro *et al.*, Phys. Rev. C **75**, 034613 (2007); K. Tsushima, H. Kim, K. Saito, Phys. Rev. C **70**, 038501 (2004)


Outline

- Electron-nucleus scattering
 - ▶ Example model: Relativistic distorted-wave impulse approximation (RDWIA)
- Nucleons in the nuclear medium
 - ▶ Example model: Quark meson coupling model (QMC)
- Experimental results
 - ▶ Polarization-transfer technique
 - ▶ $^1\text{H}(e,e'p)$: Free proton electromagnetic form factors
 - ▶ $^4\text{He}(e,e'p)$: Search for proton medium modifications
- Summary

Quasielastic Scattering from Bound Nucleons

$A(e,e'p)$: Nucleon one-body current in relativistic distorted-wave impulse approximation (RDWIA)

$$J_N^\mu(\omega, \vec{q}) = \int d\vec{p} \, \bar{\psi}_F(\vec{p} + \vec{q}) \hat{J}_N^\mu(\omega, \vec{q}) \psi_B(\vec{p})$$



- Relativistic ψ_B and ψ_F wave functions for **initial bound** and **final outgoing** nucleons, respectively.
- Relativistic **nucleon current operator** of cc1 or cc2 forms


$$\hat{J}_N^\mu(\text{cc1}) = (F_1 + \bar{\kappa}F_2)\gamma^\mu - \frac{\bar{\kappa}F_2}{2M}(P_F + \bar{P}_I)^\mu$$

(possible medium-modified form factors enter here)


Excellent Description of Many Observables

$^{16}\text{O}(e,e'p)$ at $Q^2 = 0.8 \text{ (GeV/c)}^2$

Cross section



Left-Right Asymmetry A_{LT}



- Importance of **fully relativistic calculation**; here, particularly, the bound-state spinor distortion
- Also excellent description of $^{12}\text{C}(e,e'p)$ **induced polarization**.


Nucleon in the Nuclear Medium

QCD vs. conventional Nuclear Physics

- **Conventional Nuclear Physics:**
Nucleons are effectively and well described as
 - ▶ point-like protons and neutrons (+ form factor)
 - ▶ interaction through effective forces (meson exchange)
- Underlying theory: **QCD**
 - ▶ Nucleons and mesons are not the fundamental entities
 - ▶ In the chiral limit, phase transition to quark-gluon plasma

The EMC Effect

- Depletion of the nuclear structure function $F_2^A(x)$ in the valence-quark regime $0.3 \leq x \leq 0.8$
- J. Smith and G. Miller: chiral quark-soliton model of the nucleon
Conventional nuclear physics does not explain EMC effect



- → Nucleon structure is modified in the nuclear medium

SLAC-E139 data for Iron and Gold;

Figure from Jason R. Smith and Gerald A. Miller, Phys. Rev. Lett. **91**, 212301 (2003)

Limits for Medium Modifications

- Best constraints from **y-scaling**
 - ▶ $Q^2 > 1 \text{ (GeV/c)}^2$, $\Delta G_M < 3\%$ [1]
- **Coulomb Sum Rule**, L-Response
 - ▶ No quenching in the data observed [2]
 - ▶ Quenching of S_L is experimentally established [3]
 - ▶ $Q^2 \leq 0.5 \text{ (GeV/c)}^2$: $\Delta G_E < 15\%$ or even $< 5\%$
- **Exclusive $A(e,e'p)$ processes**
 - ▶ LT Separation


[1] I. Sick, in: H. Klapdor (Ed.), Proc. Int. Conf. on Weak and Electromagnetic Interactions in Nuclei, Springer-Verlag, Berlin, 1986, p. 415

[2] J. Jourdan, Nucl. Phys. A **603**, 117 (1996), J. Carlson *et al.*, Phys. Lett. B **553**, 191 (2003)

[3] J. Morgenstern, Z.-E. Meziani, Phys. Lett. B **515**, 269 (2001)


Quark Meson Coupling Model (QMC)

- **Structure of the nucleon** described by valence quarks in a bag (Cloudy-bag model).



intermediate baryon restricted to N or Δ

- **Nuclear system** described using effective scalar (σ) and vector (ω) meson fields.





- Scalar and vector fields of nuclear matter couple directly to confined quarks.

→ Modification of **internal structure** of bound nucleon

QMC: D.H. Lu, A.W. Thomas, K. Tsushima, A.G. Williams, K. Saito, Phys. Lett. B **417**, 217 (1998) D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999); **Other models:** e.g.: J.R. Smith and G.A. Miller, Phys. Rev. C **70**, 065205 (2004); T. Horikawa, W. Bentz, Nucl. Phys. A **762**, 102 (2005)

Bound Proton EM Form Factors




- Electromagnetic rms radii and magnetic moments of the bound proton are increased
- **Charge form factor** much more sensitive to the nuclear medium than the **magnetic** ones.

D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999)

Effect on charged-current ν -A scattering

QMC bound nucleon form factors of vector and axial-vector currents



- Tsushima *et al.* compute the inclusive $^{12}\text{C}(\nu_\mu, \mu^-)X$ cross sections using a relativistic Fermi gas model with the calculated bound nucleon form factors.
- The effect of the bound nucleon form factors for this reaction is a **reduction of 8% for the total cross section**, relative to that calculated with the free nucleon form factors.

K. Tsushima, Hungchong Kim, and K. Saito, Phys. Rev. C **70**, 038501 (2004)

Polarization-Transfer Technique

- **Free** electron-nucleon scattering

$$\frac{G_E}{G_M} = -\frac{P'_x}{P'_z} \cdot \frac{(E_i + E_f)}{2m} \tan\left(\frac{\theta_e}{2}\right)$$


- **Bound** nucleons → evaluation within model

Reaction-mechanism effects in $A(\vec{e}, e' \vec{p})B$
predicted to be small and minimal for

- ▶ Quasielastic scattering
- ▶ Low missing momentum
- ▶ Symmetry about $\mathbf{p}_m = 0$

R. Arnold, C. Carlson, and F. Gross, Phys. Rev. C **23**, 363 (1981); for reaction-mechanism effects, e.g., J.M. Laget, Nucl. Phys. A **579**, 333 (1994), J.J. Kelly, Phys. Rev. C **59**, 3256 (1999), A. Meucci, C. Guisti, and F.D. Pacati, Phys. Rev. C **66**, 034610 (2002).

Proton Elastic Form-Factor Ratio




- Systematic decrease of G_E / G_M indicating difference in spatial distribution of charge and magnetization currents in the proton.

Figure from: I.A. Qattan, Phys. Rev. Lett. **94**, 142301 (2005)

Proton Charge and Magnetization Densities

Parameterization of nucleon e.m. form factors based upon radial densities



- Proton **charge density** is **broader** than magnetization density

$$\langle r_E^2 \rangle > \frac{1}{\mu_p} \langle r_M^2 \rangle$$

- Consistent with polarizabilities

$$\alpha_E > \beta_M$$

J.J. Kelly, Phys. Rev. C **66**, 065203 (2002)

S. Kondratyuk, K. Kubodera, and F. Myhrer, Phys. Rev. C **71**, 028201 (2005)

Thomas Jefferson National Accelerator Facility




JLab in Newport News, VA

- Electron-beam accelerator
- Polarized electron beam
- Beam energies up to $E_0 = 6 \text{ GeV}$
- Three experimental Halls A, B, and C

E93-049 and E03-104 at Jefferson Lab Hall A

$^4\text{He}(e, e' \vec{p})^3\text{H}$ in quasielastic kinematics $Q^2 = 0.5 - 2.6 \text{ (GeV/c)}^2$



- Polarization-transfer ratio P'_x/P'_z : sensitive to G_E/G_M
- Induced polarization P_y : sensitive to final-state interactions



S. Dieterich, *et al.*, Phys. Lett. **B500**,47(2001); S. Strauch, *et al.*, Phys. Rev. Lett. **91**, 052301(2003); JLab E03-104, R.Ent, R. Ransome, S. Strauch, P. Ulmer (spokespersons)

Polarization Measurement


Focal-Plane Polarimeter



Observed angular distribution


$$\begin{aligned}
 I(\vartheta, \varphi) &= I_0(\vartheta) (1 + \epsilon_y \cos \varphi + \epsilon_x \sin \varphi) \\
 &= I_0(\vartheta) [1 + A_C (P_y \cos \varphi - P_x \sin \varphi)]
 \end{aligned}$$

Free Proton Form-Factor Ratio G_E/G_M



- Preliminary results from E03-104 with small statistical uncertainties $\delta(P'x/P'z) \approx 0.7 \%$

$^4\text{He}(\vec{e}, e' \vec{p})$ - Polarization-Transfer Ratio




- RDWIA and RMSGA models can not describe the data.

RDWIA: J.M. Udias *et al.*, Phys. Rev. Lett. **83**, 5451 (1999);


RMSGA: P. Lava *et al.*, Phys. Rev. C **71**, 014605 (2005), D. Debruyne *et al.*, Phys. Rev. C **62**, 024611 (2000)

$^4\text{He}(\vec{e}, e' \vec{p})$ - Polarization-Transfer Ratio




- $R^{\text{RDWIA}} \approx 0.97 \times R^{\text{RPWIA}}$
- Small sensitivity to
 - bound-state wave function
 - current operator
 - optical potential
- Enhancement of lower components (spinor distortions) in RDWIA

Limited Role of MEC in ${}^4\text{He}(e,e'p){}^3\text{H}$



Relativistic mean-field calculation: A. Meucci, C. Giusti, and F.D. Pacati, Phys. Rev. C **66**, 034610 (2002)

Spin-Dependent Charge-Exchange FSI



- R suppressed by about 4% from **MEC**
- Spin-dependent **charge exchange** FSI suppresses R by about 6%
- CH-EX term not well constrained $\Rightarrow$ need P_y from E03-104


RDWIA: J.M. Udias *et al.*, Phys. Rev. Lett. **83**, 5451 (1999)

RMSGa: P. Lava *et al.*, Phys. Rev. C **71**, 014605 (2005), D. Debruyne *et al.*, Phys. Rev. C **62**, 024611 (2000)

R. Schiavilla *et al.*, Phys. Rev. Lett. **94**, 072303 (2005)

Induced Polarization in $^4\text{He}(e, e' \vec{p})$

- P_y is a measure of **final-state interactions**




R. Schiavilla *et al.*
(results **not**
acceptance averaged;
acceptance averaged
RDWIA results are
typically 40%
larger than point
values)

- Final-state interactions small, RDWIA results consistent with data
- Need smaller systematic uncertainties from **E03-104** to constrain models

In-Medium Form Factors

$$G_{E,M}(Q^2, \rho) = \frac{G_{E,M}^{\text{QMC}}(Q^2, \rho)}{G_{E,M}^{\text{QMC}}(Q^2, 0)} G_{E,M}^{\text{free}}(Q^2)$$



- Data effectively described by **proton medium modified form factors**
- New data set additional tight constraints

Summary

- **Proton in the nuclear medium**
 - ▶ Models (like **Quark Meson Coupling**) predict change of the internal structure of a bound nucleon
 - ▶ Corrections due to in-medium form factors (electromagnetic, axial) could be significant
- **Polarization transfer in $^4\text{He}(e,e'p)$**
 - ▶ Significant deviation from RDWIA results; data **effectively described by proton medium modifications**
 - ▶ Alternative interpretation in terms of strong **charge-exchange FSI**
 - ▶ Induced polarization crucial to clarify role of FSI
 - ▶ New results from E03-104 will provide needed constraints